

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 403/06, A61K 31/40, C07D 409/14, 403/14, 405/14		A1	(11) International Publication Number: WO 00/35909
			(43) International Publication Date: 22 June 2000 (22.06.00)
(21) International Application Number: PCT/EP99/09673		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 9 December 1999 (09.12.99)		<p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(30) Priority Data: 60/112,590 17 December 1998 (17.12.98) US 60/149,028 16 August 1999 (16.08.99) US			
(71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH).			
(72) Inventors: CORBETT, Wendy, Lea; 36 Ridgewood Drive, Randolph, NJ 07869-3754 (US). LUK, Kin-Chun; 66 Evergreen Drive, North Caldwell, NJ 07006-4622 (US). MAHANEY, Paige, E.; 243 Country Club Lane, Scotch Plains, NJ 07076 (US).			
(74) Agent: LOESCHNER, Thomas; Grenzacherstrasse 124, CH-4070 Basle (CH).			

(54) Title: 4-ARYLOXINDOLES AS INHIBITORS OF JNK PROTEIN KINASES



(I)

(57) Abstract

Novel 4-aryloxindoles having formula (I), where, R², R³, X and A as set forth in the specification inhibit or modulate protein kinases, in particular JNK protein kinases and are useful as anti-inflammatory agents, particularly in the treatment of rheumatoid arthritis.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

4-ARYLOXINDOLES AS INHIBITORS OF JNK PROTEIN KINASES

5

Protein kinases are a class of proteins that regulate a variety of cellular functions. This is accomplished by the phosphorylation of specific amino acids on protein substrates resulting in conformational alteration of the substrate protein. The
10 conformational change modulates the activity of the substrate or its ability to interact with other binding partners. The enzyme activity of the protein kinase refers to the rate at which the kinase adds phosphate groups to a substrate. It can be measured, for example, by determining the amount of a substrate that is converted to a product as a function of time. Phosphorylation of a substrate occurs at the active-site of a
15 protein kinase.

The JNK (Jun N-terminal kinase) protein kinases (also know as "stress-activated protein kinases" or "SAPK") are members of the mitogen-activated protein (MAP) kinases. *See, e.g.,* S. Gupta et al., EMBO J., vol. 15 no. 11 (1996) pp. 2760-
20 2770; and Yang et al., Nature, vol. 389 (23 October 1997) pp. 865-870. At least ten JNK isoforms are currently known. *See, Gupta, id.* As its name indicates, one of the substrates for JNK is c-Jun. JNK phosphorylates the NH₂-terminal activation domain of c-Jun on Ser63 and Ser73, causing increased c-Jun transcriptional activity. *See Gupta, id.* In turn, c-Jun is an AP-1 transcription factor that mediates immediate-early
25 gene expression. *See, e.g.,* A. Minden et al., Biochimica et Biophysica Acta 1333 (1997) F85-F104; and A. Karin, Biochimica et Biophysica Acta, vol.172 (1991) pp.129-157.

The JNK protein kinase is markedly activated in response to treatment of cells
30 with pro-inflammatory cytokines or exposure to environmental stress. JNK thus mediates the effect of extracellular stimuli on c-Jun. *See Gupta, supra;* and Minden,

supra. Accordingly, JNK is a physiological regulator of AP-1 transcriptional activity. Thus, inhibition of JNK activity will inhibit AP-1-dependent transcription of inflammatory and immune mediators which are implicated in pathological proliferative conditions, for example inflammatory diseases and neuro-degenerative diseases, in particular, rheumatoid arthritis. See, e.g. Swantek et al., Molecular and Cellular Biology, vol. 17 (1997) pp. 6274-6282; Maroney et al., J. Neuroscience, vol. 18 (1 Jan. 1998) pp. 104-111; and Minden, *supra*, at F92.

The rat homologue of JNK is also called SAPK (stress-activated protein kinase). SAPK isoforms share significant (>90%) sequence identity with the corresponding JNK isoforms [*compare* Kyriakis et al., Nature, Vol. 369 (12 May 1994) pp. 156-160 and Gupta et al., *supra*]. Both JNK and SAPK are capable of phosphorylation of the c-Jun substrate and thus have very similar enzyme activity. JNK and SAPK are part of a protein kinase cascade that is activated by various extracellular stimuli. See e.g. Minden *supra*; and Kyriakis et al., BioEssays Vol. 18 (1996) pp. 567-577. JNK and SAPK each can be activated by phosphorylation on specific threonine and tyrosine residues by dual specificity MAP kinase kinases such as MKK4, SEK-1, or MKK7. See Kyriakis et al., *supra*; and Tournier et al., Proceedings of the National Academy of Sciences USA Vol. 94 (July 1997), pp. 7337-7342). The dual specificity MAP kinase kinases can be activated by phosphorylation on serine and/or threonine residues by MAP kinase kinase kinases such as MEKK-1. Thus, measurement of JNK or SAPK enzyme activity may be enhanced by activation by the upstream or preceding kinases. Moreover, measurement of SAPK inhibition closely correlates with JNK inhibition.

25

Inhibitors of protein kinase catalytic activity are known in the art. See US Pat. No. 5,792,783 (3-heteroaryl-2-indolinones that modulate/inhibit tyrosine kinase signal transduction); WO 98/24432 (indoline compounds that inhibit *FLK* protein kinase); WO 97/45409 (substituted tetrahydropyrimethen-oxindole analogues that inhibit tyrosine kinase). In particular, small molecule inhibitors typically block the binding of substrates by tightly interacting with the protein kinase ATP binding site (or "active

30

site"). See WO 98/24432. It is desirable to identify small-molecule compounds that may be readily synthesized and are effective in inhibiting the catalytic activity of protein kinases, in particular of the JNK protein kinases.

5 Indolinone (also known as oxindole) compounds asserted to be useful in the regulating abnormal cell proliferation through tyrosine kinase inhibition are disclosed for example in WO 96/40116, WO 98/07695, WO 95/01349, WO 96/32380, WO 96/22976, WO 96/16964, WO 98/50356 (2-indolinone derivatives as modulators of protein kinase activity); Mohammadi et. al, Science, Vol. 276, 9 May 1997, pp. 955-
10 960. Oxindole derivatives have also been described for various other therapeutic uses: 5,206,261 (improvement of cerebral function); WO 92/07830 (peptide antagonists); EP 580 502 A1 (antioxidants).


There continues to be a need for easily synthesized, small-molecule
15 compounds effective in inhibiting JNK protein kinase and thus useful in the treatment or control of pathological proliferative conditions, for example inflammatory diseases and neuro-degenerative diseases, in particular, rheumatoid arthritis. It is thus an object of this invention to provide such compounds and compositions containing such compounds.

20

The present invention relates to 4-aryloxindoles capable of inhibiting the activity of one or more JNK protein kinases. Such compounds are useful for the treatment of inflammatory diseases and neuro-degenerative diseases. In particular, the compounds of the present invention are especially useful in the treatment or
25 control of rheumatoid arthritis.

In one embodiment, the present invention is directed to 4-aryloxindoles having the following formula:

4



1

and prodrugs and pharmaceutically active metabolites of compounds of formula I, and the pharmaceutically acceptable salts of the foregoing compounds, wherein

5

A is aryl or heteroaryl, each of which optionally substituted by one or more $-OR^4$, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-NR^6R^7$, $-CN$, $-NO_2$, $-SO_2R^4$, $-SO_2NR^6R^7$, halogen, perfluoroalkyl, lower alkyl, lower alkyl substituted by (a), halogen, cycloalkyl, and/or heterocycle; cycloalkyl or cycloalkyl substituted by (a), halogen, lower alkyl, and/or heterocycle; heterocycle or heterocycle substituted by (a), halogen, lower alkyl, and/or cycloalkyl;

10 where (a) is $-OR^4$, $-NR^6R^7$, $-COR^4$, $-COOR^4$, $-OCOR^4$, $-CONR^6R^7$, $-CN$, $-NO_2$, $-SO_2R^4$, or $-SO_2NR^6R^7$;

15 R^2 is hydrogen, $-OR^4$, $-COOR^4$, $-CONR^6R^7$, $-NR^6R^7$, halogen, $-NO_2$, $-CN$, $-SO_2NR^6R^7$, $-SO_2R^4$ perfluoroalkyl, lower alkyl, or lower alkyl substituted by $-OR^8$, $-NR^6R^7$, $-COR^4$, $-COOR^4$, and/or $-CONR^6R^7$;

20 R^3 is hydrogen, $-OR^4$, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, halogen, $-CN$, $-NR^6R^7$, perfluoroalkyl, lower alkyl, or lower alkyl substituted by $-OR^8$ and/or $-NR^6R^7$;

R^4 is hydrogen, lower alkyl or lower alkyl substituted by (b), cycloalkyl and/or heterocycle; cycloalkyl or cycloalkyl substituted (b), lower alkyl and/or heterocycle; heterocycle or heterocycle substituted by (b), lower alkyl and/or cycloalkyl;

where (b) is $-OR^5$, $-COOR^8$, $-COR^8$, $-CONR^8R^9$, $-NR^6R^7$, $-CN$, $-NO_2$, $-SO_2R^8$, or $-SO_2NR^8R^9$;

R^5 is hydrogen, $-COR^8$, $-CONR^8R^9$, lower alkyl or lower alkyl substituted by $-OR^9$, $-NR^9R^{10}$, $-N(COR^9)R^{10}$, $-COR^9$, $-CONR^9R^{10}$, and/or $-COOR^9$;

R^6 and R^7 are each independently hydrogen, $-COR^8$, $-COOR^8$, $-CONR^8R^9$, $-SO_2R^8$, $-SO_2NR^8R^9$, lower alkyl or lower alkyl substituted by cycloalkyl (or cycloalkyl substituted by (c), lower alkyl and/or heterocycle), heterocycle (or heterocycle substituted by (c), lower alkyl and/or cycloalkyl), aryl (or aryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle), or heteroaryl (or heteroaryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle); or

R^6 and R^7 are each independently cycloalkyl or cycloalkyl substituted by (c), lower alkyl and/or heterocycle; heterocycle (or heterocycle substituted by (c), lower alkyl and/or cycloalkyl), aryl (or aryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle), or heteroaryl (or heteroaryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle); where (c) is $-OR^5$, $-COOR^8$, $-COR^8$, $-CONR^8R^9$, $-CN$, $-NO_2$, $-SO_2R^8$, $-SO_2NR^8R^9$, $-NR^8R^9$;

or alternatively, $-NR^6R^7$ forms a ring having 3 to 7 atoms, said ring optionally including one or more additional hetero atoms and being optionally substituted by one or more of lower alkyl, $-OR^5$, $-COR^8$, $-COOR^8$, $CONR^8R^9$, and $-NR^5R^9$;

R^8 is hydrogen, lower alkyl (or lower alkyl substituted by cycloalkyl, heterocycle, aryl, heteroaryl, $-OR^9$, $-NR^9R^{10}$, and/or $-N(COR^9)R^{10}$), aryl (or aryl substituted by (d), lower alkyl, cycloalkyl, heterocycle, halogen and/or $-SO_2F$), heteroaryl (or heteroaryl substituted by (d), lower alkyl, cycloalkyl, heterocycle, halogen and/or $-SO_2F$), cycloalkyl (or cycloalkyl substituted by (d), lower alkyl, heterocycle and/or aryl), or

heterocycle (or heterocycle substituted by (d), lower alkyl, cycloalkyl and/or aryl); where (d) is $-OR^9$, $-COOR^9$, $-COR^9$, $-CONR^{10}R^9$, $-NR^{10}R^9$, $-CN$, $-NO_2$, $-SO_2R^9$, or $-SO_2NR^{10}R^9$;

5 R^9 and R^{10} are each independently hydrogen, lower alkyl or aryl; and

X is $=N-$ or $=CH-$.

The present invention is further directed to pharmaceutical compositions comprising a pharmaceutically effective amount of any one or more of the above-described compounds and a pharmaceutically acceptable carrier or excipient.

The present invention is also directed to the use of a compound of claim I, prodrugs of such compounds and/or salts thereof in the preparation of a medicament for treating and/or controlling inflammatory diseases and neuro-degenerative diseases, in particular, the treatment or control of rheumatoid arthritis. The present invention is also directed to intermediates useful in the preparation of the above-described 4-aryloxindoles.

20 As used herein, the following terms shall have the following definitions.

"Aryl" means an aromatic group having 5 to 10 atoms and consisting of one or 2 rings. Examples of aryl groups include phenyl and 1- or 2-naphthyl.

25 "Cycloalkyl" means a non-aromatic, partially or completely saturated cyclic aliphatic hydrocarbon group containing 3 to 8 atoms. Examples of cycloalkyl groups include cyclopropyl, cyclopentyl and cyclohexyl.

"Effective Amount" means an amount of at least one compound of Formula I or Formula II, or a pharmaceutically acceptable salt, prodrug or metabolite thereof, that inhibits the development or proliferation of (1) an inflammatory disease or response

and/or (2) a neuro-degenerative disease or response, such as for example, and not as a limitation, rheumatoid arthritis.

"Halogen" means fluorine, chlorine, bromine or iodine.

5

"Heteroaryl" groups are aromatic groups having 5 to 10 atoms, one or 2 rings, and containing one or more hetero atoms. Examples of heteroaryl groups are 2-, 3- or 4-pyridyl, tetrazolyl, oxadiazolyl, pyrazinyl, indolyl and quinolyl.

10

"Hetero atom" means an atom selected from N, O and S.

"Heterocycle" means a 3- to 10-membered non-aromatic, partially or completely saturated hydrocarbon group, such as tetrahydroquinolyl, which contains one or two rings and at least one hetero atom.

15

"IC₅₀" refers to the concentration of a particular 4-aryloxindole or 4-heteroaryloxindole required to inhibit 50% of the SAPK protein kinase catalytic activity. IC₅₀ can be measured, *inter alia*, using the assay described herein in Example 66.

20

"Lower Alkyl" denotes a straight-chain or branched saturated aliphatic hydrocarbon having 1 to 6, preferably 1 to 4, carbon atoms. Typical lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, t-butyl, 2-butyl, pentyl, hexyl and the like.

25

"Pharmaceutically acceptable salt" refers to conventional acid-addition salts or base-addition salts which retain the biological effectiveness and properties of the compounds of formula I or formula II and are formed from suitable non-toxic organic or inorganic acids or inorganic bases. Sample acid-addition salts include those derived from inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, sulfamic acid, phosphoric acid and nitric acid, and those derived from organic acids such as p-toluenesulfonic acid, salicylic acid, methanesulfonic

30

acid, oxalic acid, succinic acid, citric acid, malic acid, lactic acid, fumaric acid, and the like. Sample base-addition salts include those derived from ammonium, potassium, sodium, and quaternary ammonium hydroxide, such as for example tetramethylammonium hydroxide.

5

"Pharmaceutically acceptable," such as pharmaceutically acceptable carrier, excipient, prodrug, etc., means pharmacologically acceptable and substantially non-toxic to the subject to which the particular compound is administered.

10

"Pharmaceutically active metabolite" means a metabolic product of a compound of formula I or formula II which is pharmaceutically acceptable and effective.

15

"Prodrug" refers to a compound that may be converted under physiological conditions or by solvolysis to any of the compounds of formula I or formula II or to a pharmaceutically acceptable salt of a compound of formula I or formula II. A prodrug may be inactive when administered to a subject but is converted *in vivo* to an active compound of formula I or formula II.

20


"Substituted," as in substituted alkyl means that the substitution can occur at one or more positions and, unless otherwise indicated, that the substituents are independently selected from the specified options.

25

In a preferred embodiment of the compounds of formula I, A is aryl or heteroaryl, each of which optionally may be substituted by $-NR^6R^7$, $-OR^4$, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-SO_2R^4$, $-SO_2NR^6R^7$, lower alkyl and/or lower alkyl substituted by $-OR^5$, $-NR^6R^7$, $-COR^4$, $-COOR^4$, and/or $-CONR^6R^7$.

In another preferred embodiment, the present invention is directed to compounds having the formula:

9



II

- and the pharmaceutically acceptable salts of the foregoing compounds, wherein R^1 , $R^{1'}$ and $R^{1''}$ are each independently hydrogen, $-OR^4$, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-NR^6R^7$, $-CN$, $-NO_2$, $-SO_2R^4$, $-SO_2NR^6R^7$, halogen, perfluoroalkyl, lower alkyl (or lower alkyl substituted by (a), halogen, cycloalkyl, and/or heterocycle), cycloalkyl (or cycloalkyl substituted by (a), halogen, lower alkyl, and/or heterocycle), heterocycle (or heterocycle substituted by (a), halogen, lower alkyl, and/or cycloalkyl); where (a) is $-OR^4$, $-NR^6R^7$, $-COR^4$, $-COOR^4$, $-OCOR^4$, $-CONR^6R^7$, $-CN$, $-NO_2$, $-SO_2R^4$, or $-SO_2NR^6R^7$; and R^2 , R^3 , R^6 , R^7 and X are as defined above for formula I.

- In another preferred embodiment of the compounds of formula I and II, R^2 is hydrogen, $-OR^4$, NO_2 , perfluoroalkyl, $-NR^6R^7$, halogen, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, lower alkyl or lower alkyl substituted by $-OR^8$ and $-NR^6R^7$, $-COR^4$, $-COOR^4$, and/or $-CONR^6R^7$. Preferred perfluoroalkyls include $-CF_3$.

R^3 is preferably hydrogen, $-OR^4$, $-NR^6R^7$, lower alkyl or lower alkyl substituted $-OR^8$ and/or $-NR^6R^7$.

- R^4 is preferably hydrogen, lower alkyl or lower alkyl substituted by $-OR^5$, $-COOR^8$, $-COR^8$, $-NR^6R^7$ and/or $-CONR^8R^9$.

R^5 is preferably $-COR^8$, $-CONR^8R^9$, or lower alkyl.

R^6 and R^7 are preferably each independently hydrogen, $-\text{COR}^8$, $-\text{COOR}^8$, $-\text{CONR}^8\text{R}^9$, $-\text{SO}_2\text{R}^8$, aryl, heteroaryl, lower alkyl or lower alkyl substituted by OR^5 , and/or $-\text{NR}^8\text{R}^9$.

- 5 R^8 is preferably hydrogen, aryl, heteroaryl, lower alkyl or lower alkyl substituted by aryl, heteroaryl, $-\text{OR}^9$, $-\text{NR}^9\text{R}^{10}$, and/or $-\text{N}(\text{COR}^9)\text{R}^{10}$.

A is preferably heteroaryl, particularly indole or substituted indole.

- 10 Also preferred are compounds of formula I and II wherein X is $=\text{CH}-$.

The following are examples of preferred compounds of formula I:

- (Z)-1,3-Dihydro-4-phenyl-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (A),
15 (Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (C),
(Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride salt (D),
(Z)-1,3-Dihydro-4-(4-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-
20 one (E),
(Z)-1,3-Dihydro-4-(3-nitrophenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (F),
(Z)-1,3-Dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(3-trifluoromethylphenyl)-2H-indol-2-one (G),
25 (Z)-1,3-Dihydro-4-(4-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (H),
(Z)-1,3-Dihydro-4-(2-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (I),
(Z)-4-(2,4-Dichlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-
30 2-one (J),
(Z)-4-(4-Chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (L).

(Z)-1,3-Dihydro-4-(2-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (M),

(Z)-1,3-Dihydro-4-(1-naphthalenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (N),

5 (Z)-4-(3-Chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (P),

(Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (R),

(Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (EE),

(Z)-1,3-Dihydro-4-phenyl-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (GG),

(Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (HH),

15 (Z)-4-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid (Q),

(Z)-3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid (BB),

(Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid (II),

(Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (RR),

(Z)-4-[5-Amino-2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (SS),

25 (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid methyl ester (WW),

(Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid (XX),

(Z)-4-[2,3-Dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester trifluoroacetate salt (AAA),

30

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-hydroxybenzamide (S),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-bromobenzamide (T),

5 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-cyanobenzamide (U),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-nitrobenzamide (V),

10 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-fluorobenzamide (W),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-nitrobenzamide (X),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-methoxybenzamide (Y),

15 (Z)-4-Amino-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]cyclohexanecarboxamide (Z),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-(fluorosulfonyl)benzamide (AA),

20 (Z)-N-[2-[[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]amino]-2-oxoethyl]-4-(fluorosulfonyl)benzamide (CC),

(Z)-1,3-Dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(2-thiophenyl)-2H-indol-2-one (B),

(Z)-1,3-Dihydro-4-(2,4-dimethoxy-6-pyrimidinyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (FF),

25 (Z)-1,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (MM),

(Z)-1,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one,

30 (Z)-5-Amino-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one,

(Z)-N-[2,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (QQ),

(Z)-1,3-Dihydro-4-(4-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (TT),

5 (Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (UU),

(Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one,

10 (Z)-5-Amino-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one,

(Z)-N-[2,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (VV),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]methanesulfonamide (K),

15 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-2-thiophenesulfonamide (O),

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-4-(phenylsulfonyl)-2-thiophenesulfonamide (DD),

20 (Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (JJ),

(Z)-1,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (KK),

(Z)-N-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-4-phenyl-1H-indol-5-yl]-2-thiopheneacetamide (LL),

25 (Z)-5-Amino-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (NN),

(Z)-N-[2,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (OO),

30 (Z)-5-Amino-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-4-phenyl-2H-indol-2-one (PP),

(Z)-1,3-Dihydro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (YY),

(Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (ZZ),

5 (Z)-1,3-Dihydro-5-fluoro-4-(4-methoxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (BBB),

(Z)-1,3-Dihydro-4-(3,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (CCC),

(Z)-1,3-Dihydro-4-(2,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (DDD),

(Z)-4-(1,3-Benzodioxol-5-yl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (EEE),

(Z)-4-(3-Aminophenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (FFF),

15 (Z)-4-(3-Amino-4-methyl-phenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (GGG),

(Z)-1,3-Dihydro-5-fluoro-4-(3-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (HHH),

(Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (III),

(Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (JJJ),

2-[3-[5-Fluoro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-2,3-dihydro-1H-indol-4-yl]-phenylamino]-acetamide (KKK), and


25 (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxymethyl-3-methoxy-phenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (LLL).

The compounds disclosed herein and covered by the above formulae may exhibit tautomerism or structural isomerism. It is intended that the invention
30 encompasses any tautomeric or structural isomeric form of these compounds, or

mixtures of such forms, and is not limited to any one tautomeric or structural isomeric form utilized within the formulae drawn above.


The compounds of formula I may be prepared by processes known in the art. Suitable processes for synthesizing these compounds are provided in the examples. Generally, these compounds may be prepared according to the following synthesis scheme:

General Step 1



10

General Step 2a



Compounds 1 and 2 are either available from commercial sources or are synthesized by methods known in the art. Compounds 1 and 2 are reacted in piperidine and an appropriate solvent to yield compound 3. Compound 3 is then reacted with compound 4, which also is available from commercial sources or is synthesized by methods known in the art, to yield a compound of formula I.

Compounds of formula II may be synthesized in an analogous manner.

20

In an alternative embodiment, the present invention is directed to pharmaceutical compositions comprising at least one compound according to the invention or a prodrug thereof, or a pharmaceutically acceptable salt of a compound
5 according to the invention or a prodrug of such compound.

These pharmaceutical compositions can be administered orally, for example, in the form of tablets, coated tablets, dragees, hard or soft gelatin capsules, solutions, emulsions or suspensions. They can also be administered rectally, for example, in
10 the form of suppositories, or parenterally, for example, in the form of injection solutions.

The pharmaceutical compositions of the present invention comprising compounds of formula I or formula II, prodrugs of such compounds, or the salts
15 thereof, may be manufactured in a manner that is known in the art, *e.g.* by means of conventional mixing, encapsulating, dissolving, granulating, emulsifying, entrapping, dragee-making, or lyophilizing processes. These pharmaceutical preparations can be formulated with therapeutically inert, inorganic or organic carriers. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as such carriers
20 for tablets, coated tablets, dragees and hard gelatin capsules. Suitable carriers for soft gelatin capsules are vegetable oils, waxes, fats, semi-solid or liquid polyols. Depending on the nature of the active substance, no carriers are generally required in the case of soft gelatin capsules. Suitable carriers for the manufacture of solutions and syrups are water, polyols, saccharose, invert sugar and glucose. Suitable
25 carriers for injection are water, alcohols, polyols, glycerin, vegetable oils, phospholipids and surfactants. Suitable carriers for suppositories are natural or hardened oils, waxes, fats and semi-liquid polyols.

The pharmaceutical preparations can also contain preserving agents,
30 solubilizing agents, stabilizing agents, wetting agents, emulsifying agents, sweetening agents, coloring agents, flavoring agents, salts for varying the osmotic pressure,

buffers, coating agents or antioxidants. They can also contain other therapeutically valuable substances, including additional active ingredients other than those of formula I or formula II.


5 As mentioned above, the compounds of formula I or formula II, prodrugs thereof, and their salts, and compositions containing these compounds are useful in the treatment or control of inflammatory diseases and neuro-degenerative diseases, in particular, in the treatment or control of rheumatoid arthritis.

10 A therapeutically effective amount of a compound in accordance with this invention means an amount of compound that is effective to prevent, alleviate or ameliorate symptoms of disease of the subject being treated. Determination of a therapeutically effective amount is within the skill in the art.

15 The therapeutically effective amount or dosage of a compound according to this invention can vary within wide limits and will be adjusted to the individual requirements in each particular case. In general, in the case of oral or parental administration to adult humans weighing approximately 70 Kg, a daily dosage of about 10 mg to about 10,000 mg, preferably from about 200 mg to about 1,000 mg,
20 should be appropriate, although the upper limit may be exceeded when indicated. The daily dosage can be administered as a single dose or in divided doses, or for parental administration, it may be given as continuous infusion.


Examples

25 The compounds of the present invention may be synthesized according to known techniques, such as for example General Scheme I provided above. The following examples illustrate preferred methods for synthesizing the compounds and formulations of the present invention.


Example 1: General Synthesis Methods and Starting Materials**Method F: Preparation of carboxylic acids from the methyl esters**

- 5 The appropriate methyl ester (0.14 mmol) was dissolved in a mixture of 2 mL tetrahydrofuran and 2 mL water. Lithium hydroxide (2.8 mmol, 20 equiv.) was added, and the reaction was stirred at room temperature for 14 hours. The tetrahydrofuran was then evaporated and 10 mL water was added. The aqueous layer was then extracted with ethyl acetate (2x10 mL). These ethyl acetate extracts were discarded.
- 10 The aqueous layer was then acidified to pH = 2 with 1 N hydrochloric acid, and extracted with ethyl acetate (4x20 mL). The combined organic extracts were washed with a saturated solution of sodium chloride and were then dried over magnesium sulfate. The ethyl acetate was then evaporated and the product was recrystallized from ethanol.

15


Method L:

- To a solution of nitro compound in 10% water in methanol was added Zn dust and NH₄Cl. The mixture was heated at reflux for 6 h then filtered through Celite® (Fisher Scientific). Filtrate was concentrated *in vacuo*. The product was purified via
- 20 either flash column chromatography (SiO₂, 230-400 mesh with ethyl acetate/hexanes as solvent) or with reverse phase HPLC (using either acetonitrile / water or acetonitrile / water / trifluoroacetic acid as solvent).

Method M:


To a mixture of amino compound in THF and saturated aqueous NaHCO₃ was added a THF solution of the acid chloride dropwise. The mixture was stirred for from 3 h to 10 days at room temperature then diluted with ethyl acetate. The phases were separated and the organic solution was washed with water then dried (MgSO₄). The product was purified via either flash column chromatography (SiO₂, 230-400 mesh with ethyl acetate/hexanes as solvent) or with reverse phase HPLC (using either acetonitrile / water or acetonitrile / water / trifluoroacetic acid as solvent).

Method N: Preparation of 3-arylmethylene-substituted oxindoles via coupling with aldehyde



A solution or suspension of the appropriate oxindole (1 mmol), and excess aldehyde (1 to 2 mmol) in 2 mL of 1% piperidine in 2-propanol was heated at between 60 to 90 °C for 1 to 24 hours. Hot water (2 mL) was added. On cooling, the crystallized product was filtered off, washed with aqueous 2-propanol and dried.

Method S: Preparation of 4-aryloxindoles or 4-heteroaryloxindoles via Palladium(0)-mediated coupling




To a solution of the appropriate 4-iodooxindole (see T. Fukuyama et. al., *J. Am. Chem. Soc.* 118:7426-7427 (1996)) or 4-bromooxindole (see T. Kosuge et. al., *Chem. Pharm. Bull.* 33(4):1414-1418 (1985)) in dimethoxyethane was added an aryl or heteroaryl boronic acid, palladium catalyst, and an aqueous solution of 2M Na₂CO₃. The mixture was heated at reflux for up to 4 days. After cooling, the reaction mixture was filtered and concentrated. The product was purified via either

flash column chromatography (SiO₂, 230-400 mesh with ethyl acetate/hexanes as solvent) or with reverse phase HPLC (using either acetonitrile / water or acetonitrile / water / trifluoroacetic acid as solvent).


5 Starting Materials

Starting Material 1: (Z)-1,3-Dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one




10 A mixture of 1,3-dihydro-4-iodo-2H-indol-2-one (404.1 mg, 1.56 mmol) (*see* Fukuyama et al., *supra*) and pyrrole-2-carboxaldehyde (163.2 mg, 1.72 mmol) (Aldrich) in 2-propanol (6.2 mL) was treated with 2 drops of piperidine. The reaction mixture was heated at reflux for 24 h and then allowed to cool to 23 °C, at which time, the reaction mixture was filtered. The solid was washed several times with cold
15 distilled water and then allowed to air dry to provide pure (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (341.8 mg, 65%) as a yellow solid which was used without further purification. (MP 256 - 258 °C)

20 Starting Materials 2: (Z)-4-Bromo-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one




A mixture of 4-bromo-1,3-dihydro-2H-indol-2-one (0.2 g, 0.94 mmol) (*see* Kosuge et al., *supra*), and excess pyrrole-2-carboxaldehyde (0.11 g, 1.13 mmol) (Aldrich) in 1% piperidine in 2-propanol (2 mL) was heated at 85 °C for 2 h. Hot water
25 (2 mL) was added. On cooling, the crystallized product was filtered off, washed with aqueous 2-propanol and dried. (Yield 0.26 g, 96%)

Starting Material 3: (Z)-1,3-Dihydro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one




5 A mixture of 1,3-dihydro-4-iodo-2H-indol-2-one (0.51 g, 1.97 mmol) (see Fukuyama et al., *supra*), and excess 3-methoxy-2-pyrrolecarboxyaldehyde (0.30 g, 2.36 mmol) (prepared according to F. Bellamy et al., *J. Chem. Research (S)* (1979), 18-19; *J. Chem. Research (M)* (1979), 0101-0116) in 1% piperidine in 2-propanol (10 mL) was heated at 85 °C for 4 h. Hot water (10 mL) was added. On cooling, the
10 crystallized product was filtered off, washed with aqueous 2-propanol and dried. (Yield 0.46 g, 64%).


Starting Material 4: 1,3-Dihydro-4-iodo-5-nitro-2H-indol-2-one



15 A mixture of concentrated sulfuric acid (0.73 mL) and concentrated nitric acid (0.14 mL) was added slowly to a solution of 1,3-dihydro-4-iodo-2H-indol-2-one (0.5 g, 1.93 mmol) (see Fukuyama et al., *supra*) in concentrated sulfuric acid (6 mL) at -5 °C with stirring. The mixture was stirred for an additional 15 min at -5 °C then poured into ice. After standing for 1 h, solid was collected by filtration and washed with
20 water, and dried in a vacuum oven to give the above product. (Yield 0.46 g, 78%).


Starting Material 5: 4-Bromo-1,3-dihydro-5-nitro-2H-indol-2-one

A mixture of concentrated sulfuric acid (3.6 mL) and concentrated nitric acid
5 (0.7 mL) was added slowly to a solution of 4-bromo-1,3-dihydro-2H-indol-2-one (2 g,
9.48 mmol) (*see* Kosuge et al., *supra*) in concentrated sulfuric acid (20 mL) at -5 °C
with stirring. The mixture was stirred for an additional 1 h at -5 °C then poured into
ice. After standing for 1 h, precipitate formed was collected by filtration and washed
with water, and dried in a vacuum oven to give the above product. (Yield 2.33 g,
10 96%).

Starting Material 6: (Z)-4-Bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one

15 A mixture of 4-bromo-1,3-dihydro-5-nitro-2H-indol-2-one (0.113 g, 0.44 mmol)
(Starting Material 5 above), and excess 3-methoxy-2-pyrrolecarboxyaldehyde (66.3
mg, 0.53 mmol) (*see* Bellamy et al., *supra*) in 1% piperidine in 2-propanol (2 mL)
was heated at 85 °C for 3 h. Hot water (2 mL) was added. On cooling, the
crystallized product was filtered off, washed with aqueous 2-propanol and dried.
20 (Yield 0.136 g, 85%).


Starting Material 7: (Z)-4-Bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one



A mixture of 4-bromo-1,3-dihydro-2H-indol-2-one (100 mg, 0.47 mmol) (see
5 Kosuge et al., *supra*) and excess 3-methoxy-2-pyrrolecarboxyaldehyde (70.8 mg,
0.57 mmol) (see Bellamy et al., *supra*) in 1% piperidine in 2-propanol (1 mL) was
heated at 85 °C for 2 h. Hot water (1 mL) was added. On cooling, the crystallized
product was filtered off, washed with aqueous 2-propanol and dried. (Yield 0.13 g,
83%).

10


Starting Material 8: 2,4-Dimethoxy-6-(tributylstannyl)-pyrimidine



2,4-Dimethoxy-6-bromopyrimidine (1.73 g, 7.9 mmol) (prepared according to
B.W. Langly et al., *J. Am. Chem. Soc.* **78**:2136 (1955)) was dissolved in dry
15 tetrahydrofuran (50 mL) under argon. The solution was cooled to -100 °C with an
ethanol/liquid nitrogen bath. *n*-Butyllithium (4.74 mL, 11.8 mmol, 2.5 M solution in
hexanes) (Aldrich) was added dropwise, very slowly (dripped down the inside of the
flask in order to precool the solution), and the reaction was stirred at -100 °C for 5
min. Tributyltin chloride (Aldrich) was then added neat, and the reaction was slowly
20 allowed to warm to room temperature where it was stirred for 1 h. A saturated
solution of sodium bicarbonate (10 mL) was then added, and the tetrahydrofuran was
evaporated *in vacuo*. The product was then extracted with chloroform (3 x 50 mL),
and the combined organic extracts were dried over magnesium sulfate. The product


was purified via flash column chromatography (5% MeOH/CHCl₃) to yield 2.17 g (64%) stannane as a clear oil.

Starting Material 9: (Z)-4-Bromo-1,3-dihydro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-5-nitro-2H-indol-2-one



A mixture of 4-bromo-1,3-dihydro-5-nitro-2H-indol-2-one (1.49 g, 5.8 mmol) (Starting Material 5 above), and excess 4-methyl-5-imidazolecarboxyaldehyde (0.83 g, 7.54 mmol) (Aldrich) in 1% piperidine in 2-propanol (15 mL) was heated at 80 °C for 4 h. Hot water (15 mL) was added. On cooling, the crystallized product was filtered off, washed with aqueous 2-propanol and dried. (Yield 1.61 g, 80%).

Starting Material 10: 1,3-Dihydro-5-fluoro-4-iodo-2H-indol-2-one




A suspension of 1,3-dihydro-1-hydroxy-4-iodo-2H-indol-2-one, (2.43 g, 9 mmol) (prepared according to the procedure of A. S. Kende et al., *Synth. Commun.*, **20**(14): 2133-2138 (1990)) in dry dichloromethane (500 mL) was cooled to -25 °C under an argon atmosphere with magnetic stirring. A solution of (diethylamino)sulfur trifluoride (DAST, 1.35 mL) (Aldrich) in dry dichloromethane (40 mL) was added dropwise at a rate such that the reaction temperature did not rise above -25 °C (about 15 min.) After stirring for an additional 30 min. at -25 °C, the reaction was quenched by the addition of saturated aqueous sodium bicarbonate solution (180 mL) and allowed to warm to room temperature. The mixture was then filtered through Celite® (Fisher) and the layers separated. The aqueous layer was extracted with dichloromethane (2 X 300 mL). The dichloromethane layers were washed with

saturated aqueous sodium chloride solution (200 mL), combined, dried (magnesium sulfate) and concentrated. Residue was purified by flash chromatography on silica gel using ethyl acetate - dichloromethane (1:7, V/V) as solvent to give 1,3-dihydro-5-fluoro-4-iodo-2H-indol-2-one (Yield 1.08 g, 43%).


5

Starting Material 11: (Z)-1,3-Dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)-methylene]-2H-indol-2-one



A mixture of 1,3-dihydro-5-fluoro-4-iodo-2H-indol-2-one (0.48 g, 1.7 mmol)
10 (Starting Material 10 above), and excess 4-methyl-5-imidazolecarboxaldehyde (0.40 g, 3.6 mmol) (Aldrich) in 1% piperidine in 2-propanol (10 mL) was heated at 90 °C for 4 h. Hot water (10 mL) was added. On cooling, the crystallized product was filtered off, washed with aqueous 2-propanol and dried. Residue was purified by reverse phase chromatography using trifluoroacetic acid - acetonitrile - water as solvent to
15 give (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)-methylene]-2H-indol-2-one as trifluoroacetate salt. (Yield 0.64 g, 100%)


Starting Material 12: (Z)-1,3-Dihydro-5-fluoro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one



20


A mixture of 1,3-dihydro-5-fluoro-4-iodo-2H-indol-2-one (1.40 g, 5.05 mmol)
(see Starting Material 10), and excess 2-pyrrolecarboxyaldehyde (0.60 g, 6.3 mmol)
(Aldrich) in 1% piperidine in 2-propanol (20 mL) was heated at 85 °C for 2.25 h. Hot
water (20 mL) was added. On cooling, the crystallized product was filtered off,
25 washed with aqueous 2-propanol and dried. (Yield 1.50 g, 84%).

Starting Material 13: (Z)-1,3-Dihydro-5-fluoro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one



5 A mixture of 1,3-dihydro-5-fluoro-4-iodo-2H-indol-2-one (0.96 g, 3.47 mmol) (see Starting Material 10), and excess 3-methoxy-2-pyrrolecarboxyaldehyde (0.52 g, 4.16 mmol) (see Bellamy et al., *J. Chem. Research (S)*, 18 –19 (1979); *J. Chem. Research (M)*, 0101 –0116 (1979)), in 1% piperidine in 2-propanol (15 mL) was
10 heated at 85 °C for 3 h. Hot water (15 mL) was added. On cooling, the crystallized product was filtered off, washed with aqueous 2-propanol and dried. (Yield 1.24 g, 93%).


15 **Example 2:** (Z)-1,3-Dihydro-4-phenyl-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (A)



A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30.4 mg, 0.090 mmol) (Starting Material 1), Et₃N (38 µL, 0.271 mmol), tri-*o*-tolylphosphine (1.7 mg, 0.006 mmol) (Aldrich), Pd(OAc)₂ (0.6 mg, 0.003
20 mmol)(Aldrich), and phenylboronic acid (16.5 mg, 0.136 mmol) (Aldrich) in 362 µL of dry N,N-dimethylformamide (Fisher Scientific) was heated at 100 °C for 24 h. The reaction mixture was allowed to cool to room temperature, concentrated *in vacuo* to remove N,N-dimethylformamide, and then directly purified by flash chromatography (Merck Silica gel 60, 70-230 mesh, 10% ethyl acetate-hexanes then 25% ethyl


acetate-hexanes elution) to provide pure (Z)-1,3-dihydro-4-phenyl-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (Yield 21.9 mg, 85%) as an orange solid (mp 184 - 185 °C).

5 **Example 3:** (Z)-1,3-Dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(2-thiophenyl)-2H-indol-2-one (B)



A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (36.9 mg, 0.110 mmol) (Starting Material 1), Et₃N (46 µL, 0.329 mmol), tri-
10 tolylphosphine (2.1 mg, 0.007 mmol) (Aldrich), Pd(OAc)₂ (0.7 mg, 0.003 mmol) (Aldrich), and 2-thiopheneboronic acid (21.1 mg, 0.165 mmol) (Aldrich) in 439 µL of dry N,N-dimethylformamide was heated at 100 °C for 24 h. The reaction mixture was allowed to cool to room temperature, concentrated *in vacuo* to remove N,N-dimethylformamide, and then directly purified by flash chromatography (Merck Silica
15 gel 60, 70-230 mesh, 10% ethyl acetate-hexanes then 25% ethyl acetate-hexanes elution) to provide pure (Z)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(2-thiophenyl)-2H-indol-2-one (Yield 26.8 mg, 83%) as an orange solid (mp 213 - 214 °C).


20 **Example 4:** (Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (C)



A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (500 mg, 1.49 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (1.49


mL, 2.98 mmol), (Ph₃P)₄Pd (86 mg, 0.074 mmol) (Aldrich), and 3-aminophenylboronic acid monohydrate (253 mg, 1.63 mmol) (Lancaster) in 10 mL of a 3:1 mixture of 1,2-dimethoxyethane:distilled water was heated at 100 °C under a nitrogen atmosphere for 96 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 35% ethyl acetate-hexanes elution) to yield pure (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (410 mg, 91%) as a yellow solid (mp 92 - 94 °C) (solid to gel).

10 **Example 5:** (Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride salt (D)




A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (19.4 mg, 0.064 mmol) (Compound C from Example 4 above) in 7 mL of ethyl acetate was treated with anhydrous hydrogen chloride gas. A solid immediately precipitated. The anhydrous hydrogen chloride gas was bubbled into the reaction mixture for 4 min. The resulting solid was filtered and was allowed to air dry to provide pure hydrochloride salt (18.3 mg, 84%) as a yellow solid (mp 177 - 180 °C).

Example 6: (Z)-1,3-Dihydro-4-(4-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (E)



5 A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (100 mg, 0.298 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (342 μ L, 0.684 mmol), (Ph₃P)₄Pd (17 mg, 0.015 mmol) (Aldrich), and 4-methoxyphenylboronic acid (52 mg, 0.342 mmol) (Aldrich) in 3 mL of a 2:1 mixture of 1,2-dimethoxyethane:distilled water was heated at 100 °C under a nitrogen
10 atmosphere for 20 h. The reaction mixture was diluted with ethyl acetate, and the organic layer was washed with distilled water. The organic layer was concentrated *in vacuo* to provide a crude yellow-brown solid. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 25% ethyl acetate-hexanes elution) yielded pure (Z)-1,3-dihydro-4-(4-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (89.3 mg, 82%) as
15 a yellow solid (mp 222-223 °C).

Example 7: (Z)-1,3-Dihydro-4-(3-nitrophenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (F)




20 A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (35 mg, 0.104 mmol) (Starting Material 1), Et₃N (58 μ L, 0.415 mmol), tri-o-

tolylphosphine (6 mg, 0.020 mmol) (Aldrich), Pd(OAc)₂ (2 mg, 0.009 mmol) (Aldrich), and 3-nitrophenylboronic acid (26 mg, 0.156 mmol) (Aldrich) in 3 mL of dry N,N-dimethylformamide was heated at 100 °C for 16 h. The reaction mixture was allowed to cool to room temperature, concentrated *in vacuo* to remove N,N-

- 5 dimethylformamide, and then directly purified by flash chromatography (Merck Silica gel 60, 70-230 mesh, 10% ethyl acetate-hexanes then 25% ethyl acetate-hexanes elution) to provide pure (Z)-1,3-dihydro-4-(3-nitrophenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (24.1 mg, 70%) as an orange solid (mp 197 - 199 °C).


- 10 **Example 8: (Z)-1,3-Dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(3-trifluoromethylphenyl)-2H-indol-2-one (G)**



A suspension of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.089 mmol) (Starting Material 1), K₂CO₃ (25 mg, 0.180 mmol),

- 15 (Ph₃P)₄Pd (5 mg, 0.004 mmol) (Aldrich), and 3-(trifluoromethyl)phenylboronic acid (20 mg, 0.110 mmol) (Aldrich) in 2.5 mL of a 1.5:1 mixture of 1,2-dimethoxyethane: distilled water was heated at reflux under a nitrogen atmosphere for 14 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 25% ethyl acetate-
- 20 hexanes then 40% ethyl acetate-hexanes elution) to yield pure (Z)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(3-trifluoromethylphenyl)-2H-indol-2-one (yield 18 mg, 56%) as a yellow solid (mp 197 - 198 °C).


Example 9: (Z)-1,3-Dihydro-4-(4-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (H)



A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.089 mmol) (Starting Material 1) 2M aqueous Na₂CO₃ solution (89 μ L, 0.178 mmol), (Ph₃P)₄Pd (5 mg, 0.004 mmol) (Aldrich), and 4-methylphenylboronic acid (15 mg, 0.110 mmol) (Aldrich) in 3 mL of a 2:1 mixture of 1,2-dimethoxyethane:distilled water was heated at 100 °C under a nitrogen atmosphere for 18 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 25% ethyl acetate-hexanes elution) to provide a crude yellow solid containing product. The solid was dissolved in chloroform and precipitation by the addition of hexanes provided pure (Z)-1,3-dihydro-4-(4-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield 19 mg, 71%) as a yellow solid (mp 217-218 °C).

15

Example 10: (Z)-1,3-Dihydro-4-(2-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (I)




A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.089 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (89 μ L, 0.178 mmol), (Ph₃P)₄Pd (3 mg, 0.003 mmol) (Aldrich), and 2-methylphenylboronic acid (15 mg, 0.110 mmol) (Aldrich) in 3 mL of a 3:1 mixture of 1,2-dimethoxyethane:

20


distilled water was heated at 100 °C under a nitrogen atmosphere for 18 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 25% ethyl acetate-hexanes elution) to provide a crude yellow oil containing product. Recrystallization
5 from chloroform-pentane yielded pure (Z)-1,3-dihydro-4-(2-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield 20 mg, 75%) as yellow needle-like crystals (mp 195-197 °C).

Example 11: (Z)-4-(2,4-Dichlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (J)
10




A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (70 mg, 0.208 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (210
15 µL, 0.420 mmol), (Ph₃P)₄Pd (10 mg, 0.009 mmol) (Aldrich), and 2,4-dichlorophenylboronic acid (44 mg, 0.231 mmol) (Lancaster) in 6 mL of a 5:1 mixture of benzene:1,2-dimethoxyethane was heated at reflux under a nitrogen atmosphere for 18 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 60%
20 ethyl acetate-hexanes elution) to yield pure (Z)-4-(2,4-dichlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield 52 mg, 69%) as a yellow-brown solid (mp 185 -187 °C).

Example 12: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-(1H-pyrrol-2-yl-methylene)-1H-indol-4-yl]phenyl]methanesulfonamide (K)



5 A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (25 mg, 0.083 mmol) (from Example 4 *supra*) in 1 mL of dry pyridine (Fisher Scientific) was treated with methanesulfonyl chloride (11.4 mg, 0.100 mmol) (Aldrich). The reaction mixture was heated at 80 °C under a nitrogen atmosphere for 10 min. The reaction mixture was allowed to cool to room temperature and then
10 directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 20% ethyl acetate-hexanes elution) to yield pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-(1H-pyrrol-2-yl-methylene)-1H-indol-4-yl]phenyl]methanesulfonamide (21 mg, 67%) as a yellow solid (mp 224 - 225 °C).

15 **Example 13:** (Z)-4-(4-Chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (L)




A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (40 mg, 0.119 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (119
20 µL, 0.238 mmol), (Ph₃P)₄Pd (5 mg, 0.004 mmol) (Aldrich), and 4-chlorophenylboronic acid (21 mg, 0.134 mmol) (Aldrich) in 4 mL of a 3:1 mixture of 1,2-dimethoxyethane:distilled water was heated at reflux under a nitrogen atmosphere for

20 h. The reaction mixture was allowed to cool to room temperature and then directly purified by column chromatography. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 20% ethyl acetate-hexanes elution) provided a crude yellow solid.

Recrystallization from chloroform-pentane yielded pure (Z)-4-(4-chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (34 mg, 89%) as a yellow solid (mp 188-190 °C).

Example 14: (Z)-1,3-Dihydro-4-(2-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (M)




10

A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (50 mg, 0.149 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (149 µL, 0.298 mmol), (Ph₃P)₄Pd (5 mg, 0.004 mmol) (Aldrich), and 2-methoxyphenylboronic acid (25 mg, 0.164 mmol) (Aldrich) in 4 mL of a 2:1 mixture of 1,2-dimethoxyethane: distilled water was heated at 100 °C under a nitrogen atmosphere for 18 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 20% ethyl acetate-hexanes elution) to provide a crude yellow solid. Recrystallization from chloroform-pentane yielded pure (Z)-1,3-dihydro-4-(2-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (Yield 37 mg, 79%) as yellow crystals (mp 223 - 225 °C).

15

20


Example 15: (Z)-1,3-Dihydro-4-(1-naphthalenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (N)



A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (40 mg, 0.119 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (119 μ L, 0.238 mmol), (Ph₃P)₄Pd (3 mg, 0.003 mmol) (Aldrich), and 1-naphthaleneboronic acid (22.5 mg, 0.131 mmol) (Lancaster) in 3 mL of a 3:1 mixture of 1,2-dimethoxyethane:distilled water was heated at 100 °C under a nitrogen atmosphere for 20 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 20% ethyl acetate-hexanes then 30% ethyl acetate-hexanes elution) to yield a crude solid. Recrystallization from chloroform-pentane provided pure (Z)-1,3-dihydro-4-(1-naphthalenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield 31 mg, 78%) as yellow crystals (mp 210 - 212 °C).

15

Example 16: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-2-thiophenesulfonamide (O)




A suspension of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (27 mg, 0.090 mmol) (from Example 4 *supra*) in 1.5 mL of pyridine (Fisher Scientific) was treated dropwise with 2-thiophenesulfonyl chloride (19 mg, 0.104) (Aldrich). The reaction mixture was stirred at room temperature for 30 min and then directly purified by flash chromatography (Merck Silica gel 60, 230-400

20

mesh, 20% ethyl acetate-hexanes then 40% ethyl acetate-hexanes elution) to provide a crude yellow solid. Recrystallization from chloroform-pentane provided pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-2-thiophenesulfonamide (17 mg, 42%) as an orange solid (mp 218 - 220 °C).


5

Example 17: (Z)-4-(3-Chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (P)




10 A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.089 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (89 µL, 0.178 mmol), (Ph₃P)₄Pd (5 mg, 0.004 mmol) (Aldrich), and 3-chlorophenylboronic acid (16 mg, 0.102 mmol) (Aldrich) in 3 mL of a 2:1 mixture of 1,2-dimethoxyethane: distilled water was heated at reflux under a nitrogen atmosphere for 16 h. The
15 reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 25% ethyl acetate-hexanes elution) to provide pure (Z)-4-(3-chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (26 mg, 90%) as a yellow solid (mp 165 - 167 °C).

20 **Example 18:** (Z)-4-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid (Q)



A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.089 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (151 μ L, 0.303 mmol), (Ph₃P)₂PdCl₂ (3 mg, 0.004 mmol) (Aldrich), and 4-carboxyphenylboronic acid (18 mg, 0.108 mmol) (Lancaster) in 4.5 mL of a 2:1 mixture of 1,2-dimethoxyethane:distilled water was heated at reflux under a nitrogen atmosphere for 20 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 40% ethyl acetate-hexanes then 50% ethyl acetate-hexanes with 1% glacial acetic acid elution) to yield pure (Z)-4-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid (yield: 33 mg, 91%) as an orange solid (mp 315 - 317 °C).

Example 19: (Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (R)




15

A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (45 mg, 0.134 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (134 μ L, 0.268 mmol), (Ph₃P)₄Pd (5 mg, 0.004 mmol) (Aldrich), and 4-hydroxyphenylboronic acid (37 mg, 0.268 mmol) (prepared according to H. Gilman et al., *J. Am. Chem. Soc.* **79**:3077-3081 (1957)) in 3 mL of a 3:1 mixture of 1,2-dimethoxyethane:distilled water was heated at reflux under a nitrogen atmosphere for 15 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 35% ethyl acetate-hexanes elution) to yield pure (Z)-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield: 32 mg, 79%) as a yellow solid (mp 271 - 273 °C).

25

Example 20: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-hydroxybenzamide (S)




5 A mixture of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (59 mg, 0.196 mmol) (from Example 4 *supra*), 4-acetoxybenzoic acid (38 mg, 0.210 mmol) (Aldrich), and 1,3-dicyclohexylcarbodiimide (41 mg, 0.199) (Aldrich) in 4 mL of chloroform was heated at reflux for 22 h. The reaction mixture was diluted with hexane. The resulting yellow precipitate was collected by suction
10 filtration to provide crude carbonic acid methyl ester 4-[3-[2-oxo-3-[(1H-pyrrol-2-yl)methylene]-2,3-dihydro-1H-indol-4-yl]-phenylcarbamoyl]-phenyl ester as a yellow solid (91 mg, 98%) which was used without further purification.

A solution of the crude carbonic acid methyl ester 4-[3-[2-oxo-3-[(1H-pyrrol-2-yl)methylene]-2,3-dihydro-1H-indol-4-yl]-phenylcarbamoyl]-phenyl ester (90 mg, 0.194
15 mmol) in 2 mL of 1,4-dioxane (Aldrich) was treated with 1.5 mL of distilled water and then powdered potassium hydroxide (25 mg, 0.445 mmol). The reaction mixture was heated at 90 °C for 1 h. The reaction mixture was then allowed to cool to room temperature and directly purified by flash chromatography (Merck Silica gel 60, 230-
20 400 mesh, 45% ethyl acetate-hexanes elution) to provide a crude yellow solid containing the desired product and a 1,3-dicyclohexylcarbodiimide derivative. Most of the 1,3-dicyclohexylcarbodiimide derivative was removed by precipitation from an ethanolic solution through the slow addition of distilled water. The precipitated solid was collected by suction filtration. The filtrate was concentrated *in vacuo* and then
25 directly purified by column chromatography. Two additional rounds of flash

chromatography (Merck Silica gel 60, 230-400 mesh, 25% ethyl acetate-hexanes elution each) provided pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-hydroxybenzamide (yield: 49 mg, 60%) as a yellow solid (mp 236 - 238 °C).


5

Example 21: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-bromobenzamide (T)




10 A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride (50 mg, 0.148 mmol) (from Example 5 *supra*) and Et₃N (41 µL, 0.296 mmol) in 3 mL of chloroform was cooled to -70 °C and then treated with 3-bromobenzoyl chloride (21 µL, 0.155 mmol) (Aldrich). The reaction mixture was stirred at -70 °C for 1 hr and then stirred at room temperature for 15 h. The reaction
15 mixture was then concentrated *in vacuo*. The resulting yellow solid was suspended in water and collected by suction filtration to provide pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-bromobenzamide (yield: 70 mg, 98%) as a yellow solid (mp 251 - 253 °C).

Example 22: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-cyanobenzamide (U)



A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-
5 2H-indol-2-one hydrochloride (40 mg, 0.118 mmol) (from Example 5 *supra*) and Et₃N
(33 μ L, 0.237 mmol) in 3 mL of chloroform was treated with 3-cyanobenzoyl chloride
(22 mg, 0.130 mmol) (Aldrich), and the resulting reaction mixture was stirred at room
temperature for 15 h. The reaction mixture was concentrated *in vacuo* to remove
chloroform. The yellow residue was suspended in distilled water, and the resulting
10 solid was collected by suction filtration to provide pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-
[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-cyanobenzamide (yield: 41 mg,
81%) as a yellow solid (mp 250 - 251 °C).


Example 23: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-nitrobenzamide (V)



A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-
2H-indol-2-one hydrochloride (40 mg, 0.118 mmol) (from Example 5 *supra*) and Et₃N


(33 μ L, 0.237 mmol) in 3 mL of chloroform was cooled to -78 °C and then treated with 3-nitrobenzoyl chloride (24 mg, 0.129 mmol) (Aldrich). The reaction mixture was then stirred at room temperature under a nitrogen atmosphere for 7 h. The reaction mixture was then concentrated *in vacuo*. The resulting yellow solid was dissolved in ethanol, and re-precipitated by the addition of distilled water to yield pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-nitrobenzamide (yield: 50 mg, 94%) as a yellow solid (mp 265 - 266 °C).

Example 24: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-fluorobenzamide (W)




A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride (25 mg, 0.074 mmol) (from Example 5 *supra*) and Et₃N (21 μ L, 0.148 mmol) in 3 mL of chloroform was treated with 4-fluorobenzoyl chloride (10.5 μ L, 0.088 mmol) (Aldrich). The resulting reaction mixture was stirred at room temperature for 8 h and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 50% ethyl acetate-hexanes elution) to provide pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-fluorobenzamide (yield: 26 mg, 83%) as a yellow solid (mp 130 - 131 °C).

Example 25: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-nitrobenzamide (X)




A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride (25 mg, 0.074 mmol) (from Example 5 *supra*) and Et₃N (20 μL, 0.150 mmol) in 3 mL of chloroform was cooled to -70 °C and then treated with 4-nitrobenzoyl chloride (16 mg, 0.085 mmol) (Aldrich). The reaction mixture was stirred at -70 °C for 1 hr and then stirred at room temperature for 15 h. The reaction mixture was then concentrated *in vacuo*. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 50% ethyl acetate-hexanes elution) provided pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-nitrobenzamide (yield: 24 mg, 72%) as a yellow-brown solid (mp 157 - 159 °C).

Example 26: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-methoxybenzamide (Y)



A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride (30 mg, 0.089 mmol) (from Example 5 *supra*) and Et₃N (24 µL, 0.172 mmol) in 3 mL of chloroform was cooled to - 70 °C and then treated with 4-methoxybenzoyl chloride (17.4 mg, 0.102 mmol) (Fluka). The reaction mixture was stirred at - 70 °C for 1 hr and then stirred at room temperature for 15 h. The reaction mixture was then concentrated *in vacuo*. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 50% ethyl acetate-hexanes elution) provided pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl) methylene]-1H-indol-4-yl]-phenyl]-4-methoxybenzamide (31 mg, 80%) as a brown solid (mp 209 - 211 °C).


Example 27: (Z)-4-Amino-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl) methylene]-1H-indol-4-yl]phenyl]cyclohexanecarboxamide (Z)



A suspension of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride (53 mg, 0.157 mmol) (from Example 5 *supra*) and 4-(tert-butoxycarbonylamino)cyclohexanecarboxylic acid (53 mg, 0.204 mmol) (prepared according to K.J. Cutrona et al., *Tetrahedron Lett.* **37**(29):5045-5048 (1996)) in 4 mL of dry N,N-dimethylformamide was treated with O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU, 77 mg, 0.204 mmol) (Advance ChemTech) and N,N-diisopropylethylamine (88 µL, 0.628 mmol) (Aldrich). The reaction mixture was heated at 60 °C under a nitrogen atmosphere for 15 h. The reaction mixture was then diluted with 20 mL of a 1N aqueous hydrochloric acid solution and extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were concentrated *in vacuo*. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 45% ethyl acetate-hexanes elution) provided pure (Z)-[4-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]carbonyl]-cyclohexyl]-carbamic acid tert-butyl ester (51.4 mg, 62%) as a yellow solid.

A solution of the (Z)-[4-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenylcarbamoyl]-cyclohexyl]-carbamic acid tert-butyl ester (51.4 mg, 0.098 mmol) in 3 mL of methylene chloride was then treated with 2 mL of trifluoroacetic acid. The reaction mixture was stirred at room temperature for 1 h, and the reaction mixture was diluted with 50 mL of ethyl acetate. The organic phase was carefully washed with a saturated aqueous sodium bicarbonate solution (3 x 20 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo* to provide a yellow-brown solid. This crude solid was washed with 70% chloroform-hexanes, and the insoluble solid was collected by suction filtration to provide pure (Z)-4-amino-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]cyclohexanecarboxamide (yield: 17 mg, 40%) as a yellow solid (mp 210 - 212 °C).

Example 28: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-(fluorosulfonyl)benzamide (AA)




A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.100 mmol) (from Example 4 *supra*) and Et₃N (14 µL, 0.100 mmol) in 5 mL of chloroform was cooled to -78 °C and then treated with 4-(fluorosulfonyl)benzoyl chloride (24 mg, 0.109 mmol) (Aldrich). The reaction mixture was stirred at -78 °C under a nitrogen atmosphere for 1 hr and then heated at reflux. The reaction mixture was allowed to cool to room temperature and then diluted with chloroform. The organic phase was then washed, in sequence, as follows: with a saturated aqueous sodium bicarbonate solution, with distilled water, with a 1N aqueous hydrochloric acid solution, and finally with a saturated aqueous sodium chloride solution. The organic phase was dried over sodium sulfate, filtered, and

concentrated *in vacuo*. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 30% ethyl acetate-hexanes elution) yielded pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-(fluorosulfonyl)benzamide (yield: 25.6 mg, 53%) as a yellow solid (mp: 148 - 151 °C).

5

Example 29: (Z)-3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid (BB)




Step A: 3-Carboxyphenylboronic acid

10 A solution of 3-formylphenylboronic acid (3.35 g, 22.34 mmol) (Lancaster) in a 1:1 mixture of distilled water-acetone (64 mL) was slowly treated with potassium permanganate (6.00 g, 37.98 mmol) in small portions. The dark reaction mixture was stirred at room temperature for 1 h and then filtered through a pad of Celite®. The pad of Celite® was washed with 200 mL of distilled water then washed with 200 mL of acetone. The filtrate was concentrated *in vacuo* to a volume of approximately 200 mL. The resulting aqueous filtrate was treated with concentrated hydrochloric acid until pH = 1 and then cooled using an ice/water bath. The desired product precipitated upon cooling. The solid was filtered and washed well with cold distilled water to provide 3-carboxyphenylboronic acid as a cream-colored solid which was
15 used without further purification. (Yield 1.95 g, 53%; mp 232 - 234 °C).
20

Step B: A solution of (Z)-1,3-dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.089 mmol) (Starting Material 1), 2M aqueous Na₂CO₃ solution (150 µL, 0.300 mmol), (Ph₃P)₂PdCl₂ (4 mg, 0.006 mmol)(Aldrich), and 3-
25 carboxyphenylboronic acid (17 mg, 0.102 mmol) (from Step A above) in 4 mL of a 3:1 mixture of 1,2-dimethoxyethane:distilled water was heated at 95 °C under a nitrogen

atmosphere for 24 h. The reaction mixture was allowed to cool to room temperature and then directly purified by flash chromatography (Merck Silica gel 60, 230-400 mesh, 50% ethyl acetate-hexanes then 100% ethyl acetate with 1% glacial acetic acid elution) to provide pure (Z)-3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid (yield: 23 mg, 78%) as an orange solid (m_p 233 - 235 °C).

Example 30: (Z)-N-[2-[[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]amino]-2-oxoethyl]-4-(fluorosulfonyl)benzamide (CC)




- Step A:** A solution of glycine tert-butyl ester hydrochloride (300 mg, 1.79 mmol) (Aldrich), Et₃N (250 µL, 1.79 mmol), and 4-(fluorosulfonyl)benzoyl chloride (400 mg, 1.79 mmol) (Aldrich) in 9 mL of dry N,N-dimethylformamide was stirred at room temperature under a nitrogen atmosphere for 12 h. The reaction mixture was diluted with ethyl acetate. The organic phase was washed with distilled water, dried over sodium sulfate, filtered, and concentrated *in vacuo* to provide crude (4-fluorosulfonyl-benzoylamino)-acetic acid tert-butyl ester (541 mg, 95%) as an oil that solidified upon sitting at room temperature. The crude (4-fluorosulfonyl-benzoylamino)-acetic acid tert-butyl ester was used without further purification.
- Step B:** A solution of the crude (4-fluorosulfonyl-benzoylamino)-acetic acid tert-butyl ester (541 mg, 1.70 mmol) (from step A above) in 2 mL of methylene chloride was treated with 2 mL of trifluoroacetic acid. The reaction mixture was stirred at room temperature for 72 h. The reaction mixture was concentrated *in vacuo* to provide crude (4-fluorosulfonyl-benzoylamino)-acetic acid (407.9 mg, 92%) as a solid that was used without further purification.

Step C: A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (30 mg, 0.100 mmol) (from Example 4 *supra*), crude (4-fluorosulfonyl-benzoylamino)-acetic acid (28 mg, 0.107 mmol) (from step B above), O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU, 45 mg, 0.119 mmol) (Advance ChemTech), 1-hydroxybenzotriazole hydrate (HOBT, 16 mg, 0.118 mmol) (Aldrich), and N,N-diisopropylethylamine (52 μ L, 0.300 mmol) (Aldrich) in 3 mL of dry N,N-dimethylformamide (Fisher Scientific) was stirred at 50 °C under a nitrogen atmosphere for 7 h then stirred at room temperature under a nitrogen atmosphere for 12 h. The reaction mixture was diluted with 40 mL ethyl acetate and washed with distilled water. The organic phase was concentrated *in vacuo*. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 40% acetone-hexanes then 70% acetone-hexanes elution) provided pure (Z)-N-[2-[[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]amino]-2-oxoethyl]-4-(fluorosulfonyl)benzamide (yield: 13 mg, 24%) as a yellow solid (mp 248 - 250 °C).

15

Example 31: (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-4-(phenylsulfonyl)-2-thiophenesulfonamide (DD)




A solution of (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (60 mg, 0.200 mmol) (from Example 4 *supra*), Et₃N (28 μ L, 0.201 mmol), and 4-benzenesulfonylthiophene-2-sulfonyl chloride (64.6 mg, 0.200 mmol) (Lancaster) in 7 mL of 1,2-dimethoxyethane (Fisher Scientific) was heated at reflux for 1 h. The reaction mixture was allowed to cool and then stirred at room

20


temperature for 15 h. The reaction mixture was concentrated *in vacuo*. Flash chromatography (Merck Silica gel 60, 230-400 mesh, 45% ethyl acetate-hexanes elution) provided pure (Z)-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-4-(phenylsulfonyl)-2-thiophenesulfonamide (yield: 42 mg, 36%) as a
5 yellow solid (mp 132 - 134 °C) (solid to gel).

Example 32: (Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (EE)



10 Using Method S above, 3-aminophenyl boronic acid (0.16 g, 1.0 mmol) (Lancaster) was coupled with (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (0.3 g, 0.94 mmol) (Starting Material 7) using (Ph₃P)₄Pd (0.11g) as catalyst in aqueous 2M Na₂CO₃ (0.94 mL, 1.88 mmol) and DME (20 mL) at reflux for 1 day to produce (Z)-4-(3-aminophenyl)-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield: 0.2 g, 65%).
15


Example 33: (Z)-1,3-Dihydro-4-(2,4-dimethoxy-6-pyrimidinyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (FF)



20 A solution of 2,4-dimethoxy-6-(tributylstannyl)-pyrimidine (137 mg, 0.32 mmol) (Starting Material 8) and (Z)-4-bromo-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (80 mg, 0.30 mmol) (Starting Material 2) in 3 mL dimethylformamide and 3 mL triethylamine was degassed by bubbling argon through the solution for 15

minutes. At this time $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ (25 mg) (Aldrich) was added and the mixture heated at 70 °C for 18 h. After cooling, water (20 mL) was added and the precipitate was filtered off and dried. The product was purified via flash column chromatography (SiO_2 , 230-400 mesh with ethyl acetate/hexanes as solvent) to yield (Z)-1,3-dihydro-4-(2,4-dimethoxy-6-pyrimidinyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one. (Yield 51 mg, 49%).

Example 34: (Z)-1,3-Dihydro-4-phenyl-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (GG)




10

Using Method S above, phenyl boronic acid (86 mg, 0.70 mmol) (Aldrich) was coupled with (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (150 mg, 0.47 mmol) (Starting Material 7) using $(\text{Ph}_3\text{P})_4\text{Pd}$ (27 mg) (Aldrich) as catalyst in aqueous 2M Na_2CO_3 (0.47 mL, 0.94 mmol) and DME (10 mL) at reflux for 2 days to give (Z)-1,3-dihydro-4-phenyl-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield: 35 mg, 23%).

15

Example 35: (Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (HH)




20

In accordance with Method S above, 4-hydroxy-phenyl boronic acid (96.6 mg, 0.70 mmol) (see Gilman, *supra*) was coupled with (Z)-4-bromo-1,3-dihydro-3-[(3-


methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (150 mg, 0.47 mmol) (Starting Material 7) using $(\text{Ph}_3\text{P})_4\text{Pd}$ (27 mg) (Aldrich) as catalyst in aqueous 2M Na_2CO_3 (0.47 mL, 0.94 mmol) and DME (10 mL) at reflux for 2 days to give (Z)-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield: 60 mg, 73%, based on 70 mg recovered starting oxindole).

Example 36: (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid (II)



Using Method S above, 4-carboxy-phenyl boronic acid (27.2 mg, 0.164 mmol) (Lancaster) was coupled with (Z)-1,3-dihydro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (50 mg, 0.137 mmol) (Starting Material 3) using $(\text{Ph}_3\text{P})_4\text{Pd}$ (4.8 mg) (Aldrich) as catalyst in DME (5 mL) and solid Na_2CO_3 (51 mg, 0.48 mmol) at reflux for 18 h to give (Z)-4-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid (yield: 15 mg, 31%).


Example 37: (Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (JJ)



Using Method S above, 4-hydroxyphenyl boronic acid (85.0 mg, 0.62 mmol) (see Gilman, *supra*) was coupled with (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (150 mg, 0.41 mmol) (Starting Material


6) using DPPFPdCl₂ (17 mg) (Aldrich) as catalyst in aqueous 2M Na₂CO₃ (0.42 mL, 0.84 mmol) and DME (15 mL) at reflux for 1 day to give (Z)-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (yield: 110 mg, 52%).

Example 38: (Z)-1,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (KK)



Using Method S above, phenyl boronic acid (75.2 mg, 0.617 mmol) (Aldrich) was coupled with (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (150 mg, 0.41 mmol) (Starting Material 6) using DPPFPdCl₂ (17 mg) (Aldrich) as catalyst in aqueous 2M Na₂CO₃ (0.42 mL, 0.84 mmol) and DME (15 mL) at reflux for 18 h to give (Z)-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (yield: 120 mg, 80%).


Example 39: (Z)-N-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-4-phenyl-1H-indol-5-yl]-2-thiopheneacetamide (LL)



Using Method M above, (Z)-5-amino-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-4-phenyl-2H-indol-2-one (26.5 mg, 0.08 mmol) (from Example 43 *infra*) was acylated with 2-thiopheneacetyl chloride (25.7 mg, 0.16 mmol) (Aldrich) in saturated aqueous NaHCO₃ (0.16 mL) and THF (2 mL) at room temperature for 3 h

to give (Z)-N-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-4-phenyl-1H-indol-5-yl]-2-thiopheneacetamide (yield: 24 mg, 67%).

5 **Example 40:** (Z)-1,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (MM)



Step A: 5-Indoleboronic acid


To a suspension of sodium hydride (122 mg, 5.1 mmol) in anhydrous tetrahydrofuran (20 mL) was added 5-bromoindole (1.0 g, 5.1 mmol) (Aldrich) at 0 °C. After 15 min. of stirring at 0 °C, the reaction was cooled to -78 °C, and tert-butyllithium (10.2 mmol, 1.7M in hexane) (Aldrich) was added dropwise (a white precipitate immediately formed). After 10 min. tri-*n*-butylborate (2.75 mL, 10.2 mmol) was added dropwise. The reaction was allowed to slowly warm to room temperature, then the suspension was poured into an ice cold solution of 1N HCl. The organic phase was separated, and the aqueous phase was washed with ether. The combined organic extracts were extracted with 1N NaOH (3x 25 mL), and the combined alkaline extracts were acidified to pH 1 with 1N HCl. The product was then extracted with ether, and the combined ether layers were dried over magnesium sulfate and concentrated *in vacuo* to yield a brownish solid. The product was recrystallized from boiling water to yield 450 mg (55%) white crystals (mp > 290 °C).

Step B: (Z)-1,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one

25 A solution of 5-indoleboronic acid (36 mg, 0.22 mmol)(from Step A above), (Z)-1,3-dihydro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (54 mg, 0.15 mmol) (Starting Material 3), and sodium carbonate (52 mg, 0.49 mmol) in 5 mL


DME and 2.5 mL water was degassed for 15 min. by bubbling argon through the solution. Dichlorobis(triphenylphosphine)palladium (II) (11 mg) (Aldrich) was then added and the reaction was heated at 90 °C for 2 days. The reaction was then poured into 100 mL water and the product was extracted with ethyl acetate (4x50 mL). The combined organic layers were dried over magnesium sulfate and concentrated *in vacuo*. The product was purified via flash column chromatography (1% CH₃OH in CHCl₃) to yield 33 mg (62%) orange powder.

Example 41: (Z)-5-Amino-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (NN)




Using Method L above, (Z)-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (84 mg, 0.22 mmol) (from Example 37 *supra*) was reduced with zinc (130 mg, 2.0 mmol) and ammonium chloride (25.9 mg, 0.48 mmol) in 10% water in methanol (20 mL) at reflux for 18 h to give (Z)-5-amino-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (yield: 38 mg, 49%).

Example 42: (Z)-N-[2,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (OO)




Using Method M above, (Z)-5-amino-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (28 mg, 0.08 mmol) (from Example 41 *supra*) was acylated with 2-thiopheneacetyl chloride (25.7 mg, 0.16 mmol) (Aldrich) in saturated aqueous NaHCO₃ (0.16 mL) and THF (2 mL) at room temperature for 3 h to give (Z)-N-[2,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (yield: 16 mg, 42%).

Example 43: (Z)-5-Amino-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-4-phenyl-2H-indol-2-one (PP)




Using Method L above, (Z)-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (30 mg, 0.08 mmol) (from Example 38 *supra*) was reduced with zinc (48.9 mg, 0.75 mmol) and ammonium chloride (9.4 mg, 0.18 mmol) in 10% water in methanol (10 mL) at reflux for 1 day to give (Z)-5-amino-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-4-phenyl-2H-indol-2-one (yield: 26.5 mg, 100%).

Example 44: (Z)-N-[2,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (QQ)




Step A: (Z)-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one

55



A solution of (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (112 mg, 0.31 mmol) (Starting Material 6), 5-indoleboronic acid (71 mg, 0.44 mmol) (from Example 40, Step A, *supra*) and sodium carbonate (110 mg, 1.03 mmol) were dissolved in 10 mL DMF and 5 mL water. The solution was degassed for 30 minutes by bubbling argon through the solution. At this time, dichlorobis(triphenylphosphine) palladium(II) (11 mg) (Aldrich) was added, and the reaction was heated, under argon, at 90 °C for 2 days. Water (20 mL) was then added and the precipitate was filtered off and dried. The product was purified via flash column chromatography (SiO₂, 230-400 mesh) with 5% MeOH/CHCl₃ to give (Z)-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one as a red powder. (Yield 38 mg, 76%).

Step B: (Z)-5-Amino-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one




To a solution of (Z)-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (33 mg, 0.08 mmol) (from Step A above) in 1 mL of a 10% water in methanol solution and 0.5 mL THF was added zinc dust (145 mg, 2.21 mmol) followed by ammonium chloride (14 mg, 0.26 mmol). The reaction was heated at gentle reflux for 4 hours, at which time the reaction mixture was filtered

through a pad of Celite[®] and rinsed thoroughly with ethyl acetate. The resulting solution was diluted with 5 mL water and the product was extracted with ethyl acetate. The combined organic extracts were dried over magnesium sulfate and concentrated. The resulting powder was recrystallized from EtOAc/hex to give (Z)-5-amino-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one as a red powder. (Yield 24 mg, 80%).

Step C: (Z)-N-[2,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (QQ)

To a solution of (Z)-5-amino-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (24 mg, 0.065 mmol) (from Step B above) in 2 mL tetrahydrofuran was added 2-thiopheneacetyl chloride (24 mg, 0.15 mmol) (Aldrich) and a saturated aqueous solution of sodium bicarbonate (0.15 mL). The reaction was stirred at room temperature for 14 hours at which time the reaction was diluted with water (10 mL), and the THF was evaporated *in vacuo*. The product was then extracted with EtOAc, and the combined organic layers were dried over magnesium sulfate and concentrated *in vacuo*. The resulting powder was recrystallized from EtOAc/Hex to give product as yellow crystals. (Yield 23 mg, 72%).

Example 45: (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (RR)



Step A: 4-Methoxycarbonyl-phenyl boronic acid

A solution of (trimethylsilyl)diazomethane (3 mL, 2 M in hexanes) (Aldrich) was added to a suspension of 4-carboxyphenylboronic acid (1.0 g, 6 mmol) (Lancaster) in ether (50 mL). After stirring for 2 h at room temperature, DMF (8 mL) was added to

obtain a clear solution. An additional portion of (trimethylsilyl) diazomethane (3 mL, 2 M in hexanes) was added. After stirring for an additional 2 h, the reaction was quenched by adding acetic acid and concentrating under reduced pressure. Residue was recrystallized from water to give product. (Yield 0.81 g, 75%).


5

Step B: (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (RR)

Using Method S above, 4-methoxycarbonyl-phenyl boronic acid (29.6 mg, 0.164 mmol) (from Step A above) was coupled with (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (50 mg, 0.137 mmol) (Starting Material 6) using DPPFPdCl₂ (5.6 mg) (Aldrich) as catalyst in aqueous 2M Na₂CO₃ (0.14 mL, 0.28 mmol) and DME (5 mL) at reflux for 18 h to give (Z)-4-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (yield: 41 mg, 71%).

15

Example 46: (Z)-4-[5-Amino-2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (SS)




20

Using Method L above, (Z)-4-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (85 mg, 0.22 mmol) (from Example 45 *supra*) was reduced with zinc (130 mg, 1.97 mmol) and ammonium chloride (25.8 mg, 0.48 mmol) in 10% water in methanol (10 mL) at reflux for 2 h to yield (Z)-4-[5-amino-2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (yield: 40 mg, 51%).

25

Example 47: (Z)-1,3-Dihydro-4-(4-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (TT)



5 **Step A: 4-Indoleboronic acid**

To a suspension of sodium hydride (130 mg, 5.41 mmol) in anhydrous tetrahydrofuran (20 mL) was added 4-bromoindole (973 mg, 4.96 mmol) (see Kosuge et al. *supra*) at 0 °C. After 15 min. of stirring at 0 °C, the reaction was cooled to -78 °C, and tert-butyllithium (10.2 mmol, 1.7 M in hexane) (Aldrich) was added dropwise (a white precipitate immediately formed). After 10 min. tri-*n*-butylborate (2.75 mL, 10.2 mmol) (Aldrich) was added dropwise. The reaction was allowed to slowly warm to room temperature, then the suspension was poured into an ice cold solution of 1N HCl. The organic phase was separated, and the aqueous phase was washed with ether. The combined organic extracts were extracted with 1N NaOH (3x 25 mL), and
10 the combined alkaline extracts were acidified to pH 1 with 1N HCl. The product was then extracted with ether, and the combined ether layers were dried over magnesium sulfate and concentrated *in vacuo* to yield a brownish solid. The product was recrystallized from boiling water to yield 367 mg (46%) white crystals.


20 **Step B: (Z)-1,3-Dihydro-4-(4-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (TT)**

A solution of 4-indoleboronic acid (64 mg, 0.40 mmol) (from Step A above), (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one
25 (100 mg, 0.31 mmol) (Starting Material 7), and sodium carbonate (35 mg, 0.33 mmol) in 3 mL DME and 1 mL water was degassed for 15 min. by bubbling argon through the solution. Dichlorobis(triphenylphosphine)palladium (II) (13 mg) (Aldrich) was then added and the reaction was heated at 90 °C for 2 days. The reaction was then

poured into 100 mL water and the product was extracted with ethyl acetate (4x50 mL). The combined organic layers were dried over magnesium sulfate and concentrated *in vacuo*. The product was purified via flash column chromatography (1% CH₃OH in CHCl₃) to yield 42 mg (38%) orange powder.

5

Example 48: (Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (UU)




Step A: 6-Indoleboronic acid

10 To a suspension of sodium hydride (130 mg, 5.41 mmol) in anhydrous tetrahydrofuran (20 mL) was added 6-bromoindole (973 mg, 4.96 mmol) (prepared according to W.A. Ayer et al., *Tetrahedron* 84(14):2919-2924 (1992)) at 0 °C. After 15 min. of stirring at 0 °C, the reaction was cooled to -78 °C, and tert-butyllithium (10.2 mmol, 1.7 M in hexane) (Aldrich) was added dropwise (a white precipitate immediately formed). After 10 min. tri-*n*-butylborate (2.75 mL, 10.2 mmol) (Aldrich) was added dropwise. The reaction was allowed to warm to room temperature slowly, then the suspension was poured into an ice cold solution of 1N HCl. The organic phase was separated, and the aqueous phase was washed with ether. The combined organic extracts were extracted with 1N NaOH (3x 25 mL), and the
15 combined alkaline extracts were acidified to pH 1 with 1N HCl. The product was then
20 extracted with ether, and the combined ether layers were dried over magnesium sulfate and concentrated *in vacuo* to yield a brownish solid. The product was recrystallized from boiling water to yield 412 mg (50%) white crystals.


Step B: (Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (UU)

A solution of 6-indoleboronic acid (36 mg, 0.22 mmol) (from Step A above),
5 (Z)-1,3-dihydro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-indol-2-one (54 mg,
0.15 mmol) (Starting Material 3) and sodium carbonate (52 mg, 0.49 mmol) in 5 mL
DME and 2.5 mL water was degassed for 15 min. by bubbling argon through the
solution. Dichlorobis(triphenylphosphine)palladium (II) (10 mol%) (Aldrich) was then
added and the reaction was heated at 90 °C for 2 days. The reaction was then
10 poured into 100 mL water and the product was extracted with ethyl acetate (4x50
mL). The combined organic layers were dried over magnesium sulfate and
concentrated *in vacuo*. The resulting product, (Z)-1,3-dihydro-4-(6-indolyl)-3-[(3-
methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one), was purified via flash column
chromatography (40% EtOAc/hex) to yield 33 mg (62%) orange powder.

15

Example 49: (Z)-N-[2,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (VV)**Step A: (Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one**


20



A solution of (Z)-4-bromo-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (100 mg, 0.28 mmol) (Starting Material 6) and 6-

indoleboronic acid (48 mg, 0.30 mmol) (from Example 48, Step A) were dissolved in 2 mL DMF and 2 mL triethylamine. The solution was degassed for 30 minutes by bubbling argon through the solution. At this time dichlorobis(triphenylphosphine) palladium(II) (20 mg, 0.029) (Aldrich) was added, and the reaction was heated, under argon, at 90 °C for 2 days. Water (20 mL) was then added and the precipitate was filtered off and dried. The product was purified via flash column chromatography (SiO₂, 230-400 mesh) with 5% MeOH/CHCl₃ to yield 64 mg (57%) (Z)-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one as a red powder.

Step B: (Z)-5-amino-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one




To a solution of (Z)-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one (48 mg, 0.12 mmol) (from Step A above) in 2 mL of a 10% water in methanol solution and 0.5 mL THF was added zinc dust (70 mg, 1.07 mmol) followed by ammonium chloride (19 mg, 0.36 mmol). The reaction was heated at gentle reflux for 4 hours, at which time the reaction mixture was filtered through a pad of Celite[®] and rinsed thoroughly with ethyl acetate. The resulting solution was diluted with 5 mL water and the product was extracted with ethyl acetate. The combined organic extracts were dried over magnesium sulfate and concentrated. The resulting powder was recrystallized from EtOAc/hex to yield 25 mg (57%) of (Z)-5-amino-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one as a red powder.

Step C: (Z)-N-[2,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide (VV)


To a solution of (Z)-5-amino-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (22 mg, 0.059 mmol) (from Step B above) in 2 mL tetrahydrofuran was added 2-thiopheneacetyl chloride (20 mg, 0.12 mmol) (Aldrich) and a saturated aqueous solution of sodium bicarbonate (0.15 mL). The reaction was stirred at room temperature for 16 hours at which time the reaction was diluted with water (10 mL), and the THF was evaporated *in vacuo*. The product was then extracted with EtOAc, and the combined organic layers were dried over magnesium sulfate and concentrated *in vacuo*. The resulting powder was recrystallized from EtOAc/Hex to give product as yellow crystals. (Yield 19 mg, 66%).

Example 50: (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid methyl ester (WW)




Using Method M above, (Z)-4-[5-amino-2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester (310 mg, 0.8 mmol) (from Example 46 *supra*) was acylated with 2-thiopheneacetyl chloride (260 mg, 1.6 mmol) (Aldrich) in saturated aqueous NaHCO₃ (1.6 mL) and THF (15 mL) at room temperature for 3 h to yield (Z)-4-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid methyl ester (yield 370 mg, 90%).

Example 51: (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid (XX)




Using Method F above, (Z)-4-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid methyl ester (70 mg, 0.14 mmol) (from Example 50 *supra*) was hydrolyzed with LiOH·H₂O (20 mg, 0.48 mmol) in THF (3 mL) and water (2 mL) at room temperature for 3 days to yield (Z)-4-[2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid (yield: 20 mg, 29%).

Example 52: (Z)-1,3-Dihydro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (YY)




Using Method S above, (Z)-4-bromo-1,3-dihydro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-5-nitro-2H-indol-2-one (100 mg, 0.29 mmol) (Starting Material 9 above) was first treated with 1,1,1,3,3,3-hexamethyldisilazane (1.53 g, 9.5 mmol) (Aldrich) then coupled with phenyl boronic acid (52.4 mg, 0.43 mmol) (Aldrich) using DPPFPdCl₂ (11.7 mg) (Aldrich) as catalyst in aqueous 2M Na₂CO₃ (0.29 mL, 0.58 mmol), DMF (3 mL) and DME (3 mL) at reflux for 1 day to yield (Z)-1,3-dihydro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one (yield 9 mg, 9%).

Example 53: (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (ZZ)



Using Method S above, 4-hydroxyphenyl boronic acid (48 mg, 0.35 mmol) (see
 5 Gilman et al., *supra*) was coupled with (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11 above) using DPPFPdCl₂ (11 mg) (Aldrich) as catalyst in aqueous 2M Na₂CO₃ (0.14 mL, 0.28 mmol), DMF (3 mL) and DME (3 mL) as solvent at reflux for 3 days to yield, after reverse phase chromatography, (Z)-1,3-dihydro-5-fluoro-4-(4-
 10 hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (yield 10 mg, 22%).


Example 54: (Z)-4-[2,3-Dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester trifluoroacetate salt (AAA)
 15



Using Method S above, 4-methoxycarbonylphenyl boronic acid (36.6 mg, 0.203 mmol) (from Example 45, Step A *supra*) was coupled with (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135
 20 mmol) (Starting Material 11 above) using DPPFPdCl₂ (11 mg) (Aldrich) as catalyst in aqueous 2M Na₂CO₃ (0.14 mL, 0.28 mmol), DMF (3 mL) and DME (3 mL) at reflux for 1 day to give (Z)-4-[2,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-

2-oxo-1H-indol-4-yl]-benzoic acid methyl ester trifluoroacetate salt (yield 23 mg, 45%).


Example 55: (Z)-1,3-Dihydro-5-fluoro-4-(4-methoxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (BBB)



A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na_2CO_3 solution (0.14 mL), $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ (11 mg, 0.0135 mmol) and 4-methoxyphenylboronic acid (51.5 mg, 0.339 mmol) (Aldrich) in a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 2 days. The reaction mixture was concentrated and the crude material was purified by reverse phase HPLC to give (Z)-1,3-Dihydro-5-fluoro-4-(4-methoxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt. (Yield 18 mg, 29%).

15

Example 56: (Z)-1,3-Dihydro-4-(3,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (CCC)




A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na_2CO_3 solution (0.14 mL), $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ (11 mg, 0.0135 mmol) and 3,4-

20


dimethoxyphenylboronic acid (61.7 mg, 0.339 mmol) (Lancaster) in a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 2 days. The reaction mixture was concentrated and the crude material was purified by C18 reverse phase chromatography to give (Z)-1,3-Dihydro-4-(3,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one. (Yield 19 mg, 37%).

Example 57: (Z)-1,3-Dihydro-4-(2,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (DDD)




A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na₂CO₃ solution (0.14 mL), (Ph₃P)₂PdCl₂ (11 mg, 0.0135 mmol) and 2,4-dimethoxyphenylboronic acid (61.7 mg, 0.339 mmol) (Lancaster) in a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 2 days. The reaction mixture was concentrated and the crude material was purified by C18 reverse phase chromatography to give (Z)-1,3-Dihydro-4-(2,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one. (Yield 15 mg, 29%).

Example 58: (Z)-4-(1,3-Benzodioxol-5-yl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt (EEE)




A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na₂CO₃ solution (0.14 mL), (Ph₃P)₂PdCl₂ (11 mg, 0.0135 mmol) and 3,4-methylenedioxybenzeneboronic acid (56.3 mg, 0.339 mmol) (Lancaster) in a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 2 days. The reaction mixture was concentrated and the crude material was purified by reverse phase HPLC to yield (Z)-4-(1,3-Benzodioxol-5-yl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt. (Yield 23 mg, 36%).

10 **Example 59: (Z)-4-(3-Aminophenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (FFF)**



A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na₂CO₃ solution (0.14 mL), (Ph₃P)₂PdCl₂ (11 mg, 0.0135 mmol) and 3-aminobenzeneboronic acid (52.5 mg, 0.339 mmol) (Lancaster) in a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 4 days. The reaction mixture was concentrated and the crude material was purified by C18 reverse phase chromatography to give (Z)-4-(3-Aminophenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one. (Yield 18 mg, 40%).

Example 60: (Z)-4-(3-Amino-4-methyl-phenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (GGG)




3-Amino-4-methylphenylboronic acid was prepared by hydrogenation of 4-methyl-3-nitrophenyl-boronic acid (TCI).

A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na₂CO₃ solution (0.14 mL), (Ph₃P)₂PdCl₂ (11 mg, 0.0135 mmol) and 3-amino-4-methylphenylboronic acid (51.2 mg, 0.339 mmol) in a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 4 days. The reaction mixture was concentrated and the crude material was purified by C18 reverse phase chromatography to give (Z)-4-(3-Amino-4-methyl-phenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one. (Yield 19 mg, 40%).

15


Example 61: (Z)-1,3-Dihydro-5-fluoro-4-(3-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (HHH)



3-tert-Butyl-dimethyl-silyloxy-phenylboronic acid was prepared according to the procedure for preparing 4-tert-butyl-dimethyl-silyloxy-phenylboronic acid of: S. Yonezawa et al., Total Synthesis of Terpenin, a Novel Immunosuppressive p-Terphenyl Derivative. *J. Org. Chem.* **1998**, 63,5831-5837.

A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one (50 mg, 0.135 mmol) (Starting Material 11), 2M aqueous Na₂CO₃ solution (0.14 mL), (Ph₃P)₂PdCl₂ (11 mg, 0.0135 mmol) and 3-tert-butyl-dimethyl-silyloxy-phenylboronic acid (0.14 g, 0.54 mmol) in 5 ml of a 1:4 mixture of DMF:1,2-dimethoxyethane (5 mL) was heated at 104 °C for 3 days. The reaction mixture was concentrated and the crude material was washed with methanol to give (Z)-1,3-Dihydro-5-fluoro-4-(3-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one. (Yield 10 mg, 22%).


10 **Example 62:** (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (III)



4-Hydroxyphenylboronic acid was prepared according to the procedure of H. Gilman et al., Hydroxybenzeneboronic Acids and Anhydrides. *J. Am. Chem. Soc.* 1957, 79, 3077-3081.


A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one (100 mg, 0.28 mmol) (Starting Material 12), 2M aqueous Na₂CO₃ solution (0.28 mL), (Ph₃P)₂PdCl₂ (22.9 mg, 0.028 mmol) and 4-hydroxyphenylboronic acid (77.2 mg, 0.56 mmol) in 1,2-dimethoxyethane (10 mL) was heated at 104 °C for 2 days. The reaction mixture was filtered and concentrated. The crude material was purified by reverse phase HPLC to give (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one. (Yield 55mg, 61%).

Example 63: (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (JJJ)



A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (100 mg, 0.26 mmol) (Starting Material 13), 2M aqueous Na₂CO₃ solution (0.26 mL), (Ph₃P)₂PdCl₂ (21.2 mg, 0.026 mmol) and 4-hydroxyphenylboronic acid (53.9 mg, 0.39 mmol) in 1,2-dimethoxyethane (10 mL) was heated at 104 °C for 4 days. The reaction mixture was filtered and concentrated. The crude material was purified by reverse phase HPLC to give (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one. (Yield 41mg, 45%).

Example 64: 2-[3-[5-Fluoro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-2,3-dihydro-1H-indol-4-yl]-phenylamino]-acetamide (KKK)




3-(Carbamoylmethyl-amino)-phenylboronic acid was prepared according to the procedure of A. H. Soloway et al., Acylation and alkylation of aminoboronic acid. *J. Org. Chem.* 1960, 25, 1683-1686.

A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (50 mg, 0.13 mmol), 2M aqueous Na₂CO₃ solution (0.13 mL) (Starting Material 13), (Ph₃P)₂PdCl₂ (22 mg, 0.027 mmol) and 3-

(carbamoylmethyl-amino)-phenylboronic acid (55 mg, 0.26 mmol) in 1,2-dimethoxyethane (5 mL) was heated at 103 °C for 3 days. The reaction mixture was filtered and concentrated. The crude material was purified by C18 reverse phase chromatography to give 2-[3-[5-Fluoro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-2,3-dihydro-1H-indol-4-yl]-phenylamino]-acetamide. (Yield 15mg, 28%).

Example 65: (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxymethyl-3-methoxy-phenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (LLL)



4-tert-Butyl-dimethyl-silyloxymethyl-3-methoxyphenylboronic acid was prepared according to the procedure of S. Yonezawa et al., supra.

A solution of (Z)-1,3-dihydro-5-fluoro-4-iodo-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one (50 mg, 0.13 mmol) (Starting Material 13), 2M aqueous Na₂CO₃ solution (0.13 mL), (Ph₃P)₂PdCl₂ (11 mg, 0.013 mmol) and 4-tert-butyl-dimethyl-silyloxymethyl-3-methoxyphenylboronic acid (77 mg, 0.26 mmol) in 1,2-dimethoxyethane (5 mL) was heated at 104 °C for 1.5 days. The reaction mixture was filtered and concentrated. The crude material was purified by chromatography with silica gel eluting with EtOAc/Hexanes (3:7) to give silylated product (50 mg) which was deprotected by treatment with tetrabutyl ammonium fluoride to give (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxymethyl-3-methoxy-phenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one. (Yield 30mg, 76%).

Example 66: SAPK Inhibitory Activity

The SAPK inhibitory activity of the compounds of the invention is demonstrated below. These effects indicate that the compounds of the present invention are useful in treating inflammatory diseases such as, for example, rheumatoid arthritis.

SAPK FlashPlate Assay

Human JNK is highly homologous to rat SAPK. To measure the inhibitory activity of test compounds, the compounds were tested in the rat SAPK assay.

For the SAPK assay, purified GST-c-Jun (a chimeric protein containing c-Jun, a natural substrate of JNK) was coated on 96 well FlashPlates (New England Nuclear, Boston, MA). Purified rat SAPK (isoform b, Kyriakis et al. *supra*) was preincubated with preparations containing MEKK-1 and MKK4 for 30 minutes at 37°C in assay buffer containing 25 mM HEPES, pH 7.5, 150 mM NaCl, 20 mM MgCl₂, 2 mM DTT, 0.001% Tween 20, 1 mM ATP freshly added. In the preincubation step, MEKK-1 phosphorylates and activates MKK-4, which in turn phosphorylates and activates SAPK. The activated SAPK was then added to the c-Jun coated FlashPlates along with ³³P-ATP (0.32 mCi per reaction) and test compounds. The plates were incubated for 30 minutes at 37°C, then washed with PBS, 0.01% Tween 20, and counted in the Topcount scintillation counter (Packard Instrument Co., Downers Grove, IL). Dilutions of compounds were tested in duplicate in each assay. The percent inhibition of c-Jun phosphorylation (a measure of inhibition of SAPK activity) was determined by the following formula:

$$100 \times \left[1 - \frac{\text{test compound} - \text{nonspecific}}{\text{total} - \text{nonspecific}} \right]$$

where "test compound" refers to the average counts per minute of the test duplicates, "nonspecific" refers to the average counts per minute when no SAPK was added, and "total" refers to the average counts per minute when no compound was added.

The results of the SAPK assay with various test compounds is summarized below in Table I.

Table I

	IC ₅₀ (μM)
Compound	SAPK
C	<0.15
EE	<0.15
GG	<0.15
JJ	<0.15
KK	<0.15
LL	<0.15
MM	<0.15
NN	<0.15
OO	<0.15
PP	<0.15
QQ	<0.15
RR	<0.15
SS	<0.15
TT	<0.15
UU	<0.15
VV	<0.15
WW	<0.15
XX	<0.15

5 U937 Cell-Based Assay

The U937 cells, a human monocyte/macrophage cell line, was obtained from the ATTC and grown in the recommended medium. These cells when stimulated with lipopolysaccharide (LPS) release TNF, another inflammatory mediator implicated in the JNK pathway (Swantek et al., *supra*) and IL-6. In this assay the ability of a test compound to block TNF expression is evaluated.

The U937 cells were suspension cells which when stimulated with phorbol myristate acetate (PMA) (Sigma, St. Louis, MO) became adherent. After PMA stimulation the cells were washed in cell culture medium and plated at 1X10⁵ cells/well in 96 well plates. The following day the test compounds and dexamethasone control (Sigma, St. Louis, MO) were added to the cells for 1 hour of preincubation. Then the cells were stimulated with LPS (Sigma, St. Louis, MO). After

an additional 24 hours of incubation the supernatants were removed and assayed for TNF- α and IL-6 by ELISA. The IL-6 ELISA was run as described previously for the MG63 assay. The TNF ELISA was run using a kit supplied by Genzyme (Cambridge, MA).

5

In the ELISA, 96 well plates were coated with antibody to TNF- α or IL-6. Supernatants were added to the coated plates and any antigen (TNF- α or IL-6) in the supernatant bound to the antibody coated on the plates. The plates were then washed with PBS containing 0.05% Tween 20 (Sigma, St. Louis, MO) and the
10 biotinylated secondary antibody was added. This secondary antibody binds to the already bound antigen creating a "sandwich effect". Plates were washed as described above and horseradish peroxidase (HRP)-streptavidin conjugate (Sigma, St. Louis, MO) was added to the plates. HRP-streptavidin bound to the biotin-antibody conjugate. The plates were washed and TMB substrate (Kirkegaard and Perry Labs,
15 Gaithersburg, MD) was added to the wells. This substrate changes color in the presence of HRP-streptavidin. The intensity of the color (measured at 450 nm) is proportional to the amount of TNF- α or IL-6 produced by the U937 cells upon exposure to LPS and the test compounds. Optical density values were converted to concentration (pg/ml or Units/ml) based on a standard curve included in the assay.
20 IC₅₀ values for each test compound were determined from the linear regression of a plot of the logarithm of the concentration of compound versus amount of TNF- α or IL-6 secreted. (The TNF- α antibodies and the IL-6 antibodies were obtained from either Genzyme, Cambridge, MA or Pharmingen, San Diego, CA.).

25

The results of this assay on various test compounds is summarized below in Table II.

Table II

Compound	IC50 (μ M) in U837 Cells	
	TNF	IL6
D	1.17	6.33
WW	1.35	7.80

5 **Example 67: Tablet Formulation**

Item	Ingredients	mg/Tablet					
		5	25	100	250	500	750
1	Compound 1*	5	25	100	250	500	750
2	Anhydrous Lactose	103	83	35	19	38	57
3	Croscarmellose Sodium	6	6	8	16	32	48
4	Povidone K30	5	5	6	12	24	36
5	Magnesium Stearate	1	1	1	3	6	9
	Total Weight	120	120	150	300	600	900

*Compound 1 represents a compound of the invention.

Manufacturing Procedure:

- 10 1. Mix Items 1, 2 and 3 in a suitable mixer for 15 minutes.
2. Granulate the powder mix from Step 1 with 20% Povidone K30 Solution (Item 4).
3. Dry the granulation from Step 2 at 50°C.
4. Pass the granulation from Step 3 through a suitable milling equipment.
5. Add the Item 5 to the milled granulation Step 4 and mix for 3 minutes.
- 15 6. Compress the granulation from Step 5 on a suitable press.

Example 68: Capsule Formulation

Item	Ingredients	mg/Capsule				
1	Compound 1 *	5	25	100	250	500
2	Anhydrous Lactose	159	123	148	--	--
3	Corn Starch	25	35	40	35	70
4	Talc	10	15	10	12	24
5	Magnesium Stearate	1	2	2	3	6
	Total Fill Weight	200	200	300	300	600

* Compound 1 represents a compound of the invention.

5 **Manufacturing Procedure:**

1. Mix Items 1, 2 and 3 in a suitable mixer for 15 minutes.
2. Add Items 4 & 5 and mix for 3 minutes.
3. Fill into a suitable capsule.

10 **Example 69: Injection Solution/Emulsion Preparation**

Item	Ingredient	mg/mL
1	Compound 1 *	1 mg
2	PEG 400	10-50 mg
3	Lecithin	20-50 mg
4	Soy Oil	1-5 mg
5	Glycerol	8-12 mg
6	Water q.s.	1 mL

* Compound 1 represents a compound of the invention.

Manufacturing Procedure:

- 15
1. Dissolve item 1 in item 2
 2. Add items 3, 4 and 5 to item 6 and mix until dispersed, then homogenize.

3. Add the solution from step 1 to the mixture from step 2 and homogenize until the dispersion is translucent.
4. Sterile filter through a 0.2 um filter and fill into vials.

5 **Example 70: Injection Solution/Emulsion Preparation**

Item	Ingredient	mg/mL
1	Compound 1 *	1 mg
2	Glycofurol	10-50 mg
3	Lecithin	20-50 mg
4	Soy Oil	1-5 mg
5	Glycerol	8-12 mg
6	Water	q.s. 1 mL

* Compound 1 represents a compound of the invention.

Manufacturing Procedure:

- 10 1. Dissolve item 1 in item 2
2. Add items 3, 4 and 5 to item 6 and mix until dispersed, then homogenize.
3. Add the solution from step 1 to the mixture from step 2 and homogenize until the dispersion is translucent.
4. Sterile filter through a 0.2 um filter and fill into vials.

15


While the invention has been illustrated by reference to specific and preferred embodiments, those skilled in the art will understand that variations and modifications may be made through routine experimentation and practice of the invention. Thus, the invention is intended not to be limited by the foregoing description, but to be

20 defined by the appended claims and their equivalents.

CLAIMS

What Is Claimed Is:

- 5 1. A compound of the formula



and the pharmaceutically acceptable salts of the foregoing compounds,
wherein

- 10 A is aryl or heteroaryl, each of which optionally substituted by one or more -OR⁴, -COR⁴, -COOR⁴, -CONR⁶R⁷, -NR⁶R⁷, -CN, -NO₂, -SO₂R⁴, -SO₂NR⁶R⁷, halogen, perfluoroalkyl, lower alkyl, lower alkyl substituted by (a), halogen, cycloalkyl, and/or heterocycle; cycloalkyl or cycloalkyl substituted by (a), halogen, lower alkyl, and/or heterocycle; heterocycle or heterocycle substituted by (a), halogen, lower alkyl,
15 and/or cycloalkyl;

where (a) is -OR⁴, -NR⁶R⁷, -COR⁴, -COOR⁴, -OCOR⁴, -CONR⁶R⁷, -CN, -NO₂, -SO₂R⁴, or -SO₂NR⁶R⁷;

- R² is hydrogen, -OR⁴, -COOR⁴, -CONR⁶R⁷, -NR⁶R⁷, halogen, -NO₂, -CN, -SO₂NR⁶R⁷,
20 -SO₂R⁴ perfluoroalkyl, lower alkyl, or lower alkyl substituted by -OR⁸, -NR⁶R⁷, -COR⁴, -COOR⁴, and/or -CONR⁶R⁷;

R³ is hydrogen, -OR⁴, -COR⁴, -COOR⁴, -CONR⁶R⁷, halogen, -CN, -NR⁶R⁷, perfluoroalkyl, lower alkyl, or lower alkyl substituted by -OR⁸ and/or -NR⁶R⁷;

R^4 is hydrogen, lower alkyl or lower alkyl substituted by (b), cycloalkyl and/or heterocycle; cycloalkyl or cycloalkyl substituted (b), lower alkyl and/or heterocycle; heterocycle or heterocycle substituted by (b), lower alkyl and/or cycloalkyl;
 5 where (b) is $-OR^5$, $-COOR^8$, $-COR^8$, $-CONR^8R^9$, $-NR^6R^7$, $-CN$, $-NO_2$, $-SO_2R^8$, or $-SO_2NR^8R^9$;

R^5 is hydrogen, $-COR^8$, $-CONR^8R^9$, lower alkyl or lower alkyl substituted by $-OR^9$, $-NR^9R^{10}$, $-N(COR^9)R^{10}$, $-COR^9$, $-CONR^9R^{10}$, and/or $-COOR^9$;

10

R^6 and R^7 are each independently hydrogen, $-COR^8$, $-COOR^8$, $-CONR^8R^9$, $-SO_2R^8$, $-SO_2NR^8R^9$, lower alkyl or lower alkyl substituted by cycloalkyl (or cycloalkyl substituted by (c), lower alkyl and/or heterocycle), heterocycle (or heterocycle substituted by (c), lower alkyl and/or cycloalkyl), aryl (or aryl substituted by (c), lower
 15 alkyl, cycloalkyl and/or heterocycle), or heteroaryl (or heteroaryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle); or

R^6 and R^7 are each independently cycloalkyl or cycloalkyl substituted by (c), lower alkyl and/or heterocycle;

heterocycle (or heterocycle substituted by (c), lower alkyl and/or cycloalkyl),
 20 aryl (or aryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle), or heteroaryl (or heteroaryl substituted by (c), lower alkyl, cycloalkyl and/or heterocycle);
 where (c) is $-OR^5$, $-COOR^8$, $-COR^8$, $-CONR^8R^9$, $-CN$, $-NO_2$, $-SO_2R^8$, $-SO_2NR^8R^9$, $-NR^8R^9$;

25 or alternatively, $-NR^6R^7$ forms a ring having 3 to 7 atoms, said ring optionally including one or more additional hetero atoms and being optionally substituted by one or more of lower alkyl, $-OR^5$, $-COR^8$, $-COOR^8$, $CONR^8R^9$, and $-NR^5R^9$;

R^8 is hydrogen, lower alkyl (or lower alkyl substituted by cycloalkyl, heterocycle, aryl,
 30 heteroaryl, $-OR^9$, $-NR^9R^{10}$, and/or $-N(COR^9)R^{10}$),


- aryl (or aryl substituted by (d), lower alkyl, cycloalkyl, heterocycle, halogen and /or –SO₂F),
heteroaryl (or heteroaryl substituted by (d), lower alkyl, cycloalkyl, heterocycle, halogen and /or –SO₂F),
5 cycloalkyl (or cycloalkyl substituted by (d), lower alkyl, heterocycle and/or aryl), or heterocycle (or heterocycle substituted by (d), lower alkyl, cycloalkyl and/or aryl);
where (d) is –OR⁹, –COOR⁹, –COR⁹, –CONR¹⁰R⁹, –NR¹⁰R⁹, –CN, –NO₂, –SO₂R⁹, or –SO₂NR¹⁰R⁹;
10 R⁹ and R¹⁰ are each independently hydrogen, lower alkyl or aryl; and
X is =N- or =CH-.

2. The compounds of claim 1, wherein
15 R⁸ is hydrogen, lower alkyl optionally substituted by cycloalkyl, heterocycle, aryl, heteroaryl, –OR⁹, –NR⁹R¹⁰, and/or –N(COR⁹)R¹⁰;
aryl optionally substituted by the group consisting of (d), lower alkyl, cycloalkyl and/or heterocycle;
heteroaryl optionally substituted by (d), lower alkyl, cycloalkyl and/or heterocycle;
20 cycloalkyl optionally substituted by (d), lower alkyl, heterocycle and/or aryl;
heterocycle optionally substituted by (d), lower alkyl, cycloalkyl and/or aryl;
where (d) is –OR⁹, –COOR⁹, –COR⁹, –CONR¹⁰R⁹, –NR¹⁰R⁹, –CN, –NO₂, –SO₂R⁹, or –SO₂NR¹⁰R⁹;
R⁹ and R¹⁰ are each independently hydrogen or lower alkyl;
25 and R², R³, A and X are as in claim 1.

3. The compounds of claim 1 or 2 wherein A is aryl or heteroaryl each of which optionally is substituted by –NR⁶R⁷, –OR⁴, –COR⁴, –COOR⁴, –CONR⁶R⁷, –SO₂R⁴, –SO₂NR⁶R⁷, lower alkyl and/or lower alkyl substituted by –OR⁵, –NR⁶R⁷, –COR⁴, –COOR⁴, and/or –CONR⁶R⁷.
30

81

4. The compounds of claim 1 having the formula:



and the pharmaceutically acceptable salts thereof wherein

- 5 R¹, R¹' and R¹'' are each independently hydrogen, -OR⁴, -COR⁴, -COOR⁴, -CONR⁶R⁷, -NR⁶R⁷, -CN, -NO₂, -SO₂R⁴, -SO₂NR⁶R⁷, halogen, perfluoroalkyl, lower alkyl (or lower alkyl substituted by (a), halogen, cycloalkyl, and/or heterocycle), cycloalkyl (or cycloalkyl substituted by (a), halogen, lower alkyl, and/or heterocycle), heterocycle (or heterocycle substituted by (a), halogen, lower alkyl, and/or cycloalkyl);
- 10 where (a) is -OR⁴, -NR⁶R⁷, -COR⁴, -COOR⁴, -OCOR⁴, -CONR⁶R⁷, -CN, -NO₂, -SO₂R⁴, or -SO₂NR⁶R⁷;
- and R², R³, R⁶, R⁷ and X are as defined in claim 1 for formula I.

5. The compounds of any one of claims 1 to 4 wherein R² is hydrogen, -OR⁴, -NO₂, -NR⁶R⁷, perfluoroalkyl, halogen, -COR⁴, -COOR⁴, -CONR⁶R⁷ lower alkyl and/or lower alkyl substituted by -OR⁸ and/or -NR⁶R⁷.
- 15

6. The compounds of any one of claims 1 to 5 wherein R³ is hydrogen, -OR⁴, -NR⁶R⁷, lower alkyl and/or lower alkyl substituted by the group consisting of -OR⁸ and -NR⁶R⁷.
- 20

7. The compounds of any one of claims 1 to 6 wherein R⁴ is selected from the group consisting of -H and lower alkyl which optionally may be substituted by the group consisting of -OR⁵, -COOR⁸, -COR⁸, -NR⁶R⁷ and -CONR⁸R⁹.

8. The compounds of any one of claims 1 to 7 wherein R^5 is selected from the group consisting of $-\text{COR}^8$, $-\text{CONR}^8\text{R}^9$, and lower alkyl.
- 5 9. The compounds of any one of claims 1 to 8 wherein R^6 and R^7 are each independently selected from the group consisting of $-\text{H}$, $-\text{COR}^8$, $-\text{COOR}^8$, $-\text{CONR}^8\text{R}^9$, $-\text{SO}_2\text{R}^8$, aryl, heteroaryl and lower alkyl which optionally may be substituted by the group consisting of OR^5 , and $-\text{NR}^8\text{R}^9$.
- 10 10. The compounds of any one of claims 1 to 9 wherein R^8 is selected from the group consisting of $-\text{H}$, aryl, heteroaryl and lower alkyl which optionally may be substituted by the group consisting of aryl, heteroaryl, $-\text{OR}^9$, $-\text{NR}^9\text{R}^{10}$, and $-\text{N}(\text{COR}^9)\text{R}^{10}$.
- 15 11. The compounds of any one of claims 1 to 10 wherein X is $=\text{CH}-$.
12. The compound of claim 1 or 2 wherein A is heteroaryl.
- 20 13. The compound of claim 12 wherein A is indole or substituted indole.
14. A compound of claim 5 which is
- 25 (Z)-1,3-Dihydro-4-phenyl-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;
(Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;
(Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one hydrochloride salt;
- 30 (Z)-1,3-Dihydro-4-(4-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;
(Z)-1,3-Dihydro-4-(3-nitrophenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;
(Z)-1,3-Dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(3-trifluoromethylphenyl)-2H-indol-2-one;

(Z)-1,3-Dihydro-4-(4-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one ;

(Z)-1,3-Dihydro-4-(2-methylphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

5 (Z)-4-(2,4-Dichlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-4-(4-Chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

10 (Z)-1,3-Dihydro-4-(2-methoxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-1,3-Dihydro-4-(1-naphthalenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-4-(3-Chlorophenyl)-1,3-dihydro-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

15 (Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-4-(3-Aminophenyl)-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

20 (Z)-1,3-Dihydro-4-phenyl-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one; or

(Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one.

15. A compound of claim 5 which is

25 (Z)-4-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid;

(Z)-3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-benzoic acid;

30 (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid;

(Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester;

(Z)-4-[5-Amino-2,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester;

5 (Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid methyl ester;

(Z)-4-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-5-[(2-thienylacetyl)amino]-1H-indol-4-yl]-benzoic acid;

10 (Z)-4-[2,3-Dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2-oxo-1H-indol-4-yl]-benzoic acid methyl ester trifluoroacetate salt;

16. A compound of claim 5 which is

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-hydroxybenzamide;

15 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-bromobenzamide;

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-cyanobenzamide;

20 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-3-nitrobenzamide;

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-fluorobenzamide;

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-nitrobenzamide;

25 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-methoxybenzamide;

(Z)-4-Amino-N-[3-[2,3-dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]cyclohexanecarboxamide;

30 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]-phenyl]-4-(fluorosulfonyl)benzamide; or

(Z)-N-[2-[[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]amino]-2-oxoethyl]-4-(fluorosulfonyl)benzamide;

17. A compound of claim 12 which is

5 (Z)-1,3-Dihydro-3-[(1H-pyrrol-2-yl)methylene]-4-(2-thiophenyl)-2H-indol-2-one;

(Z)-1,3-Dihydro-4-(2,4-dimethoxy-6-pyrimidinyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-1,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

10 (Z)-1,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one,

(Z)-5-Amino-1,3-dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one,

15 (Z)-N-[2,3-Dihydro-4-(5-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide;

(Z)-1,3-Dihydro-4-(4-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

20 (Z)-1,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one,

(Z)-5-Amino-1,3-dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one, or

25 (Z)-N-[2,3-Dihydro-4-(6-indolyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide.

18. A compound of claim 5 which is

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]methanesulfonamide;

30 (Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-2-thiophenesulfonamide; or

(Z)-N-[3-[2,3-Dihydro-2-oxo-3-[(1H-pyrrol-2-yl)methylene]-1H-indol-4-yl]phenyl]-4-(phenylsulfonyl)-2-thiophenesulfonamide.

19. A compound of claim 5 which is

5 (Z)-1,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-2H-indol-2-one;

(Z)-1,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one;

10 (Z)-N-[2,3-Dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-4-phenyl-1H-indol-5-yl]-2-thiopheneacetamide;

(Z)-5-Amino-1,3-dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

(Z)-N-[2,3-Dihydro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-1H-indol-5-yl]-2-thiopheneacetamide;

15 (Z)-5-Amino-1,3-dihydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-4-phenyl-2H-indol-2-one;

(Z)-1,3-Dihydro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-5-nitro-4-phenyl-2H-indol-2-one; or

20 (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt.

20. A compound of claim 1 which is

25 (Z)-1,3-Dihydro-5-fluoro-4-(4-methoxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt;

(Z)-1,3-Dihydro-4-(3,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one;

(Z)-1,3-Dihydro-4-(2,4-dimethoxyphenyl)-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one;

30 (Z)-4-(1,3-Benzodioxol-5-yl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one trifluoroacetate salt;

(Z)-4-(3-Aminophenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one;

(Z)-4-(3-Amino-4-methyl-phenyl)-1,3-dihydro-5-fluoro-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one;

5 (Z)-1,3-Dihydro-5-fluoro-4-(3-hydroxyphenyl)-3-[(4-methyl-1H-imidazol-5-yl)methylene]-2H-indol-2-one;

(Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

10 (Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxyphenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one;

2-[3-[5-Fluoro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-oxo-2,3-dihydro-1H-indol-4-yl]-phenylamino]-acetamide; or

(Z)-1,3-Dihydro-5-fluoro-4-(4-hydroxymethyl-3-methoxy-phenyl)-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-indol-2-one.

15

21. (Z)-1,3-Dihydro-4-iodo-3-[(1H-pyrrol-2-yl)methylene]-2H-indol-2-one

22. A pharmaceutical composition comprising as an active ingredient a compound of claim 1 and a pharmaceutically acceptable carrier or excipient.

20

23. The compounds of claim 1 for use as medicaments.

24. The use of a compound of claim 1 in the preparation of a medicament containing a compound of claim 1 for treating a neuro-degenerative disease,
25 particularly rheumatoid arthritis.

25. The novel compounds, compositions and use as hereinbefore described, especially with reference to the Examples.

30

INTERNATIONAL SEARCH REPORT

In **International Application No**
PCT/EP 99/09673

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D403/06 A61K31/40 C07D409/14 C07D403/14 C07D405/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 40116 A (SUGEN, INC.) 19 December 1996 (1996-12-19) cited in the application page 1 -page 2; claim 1 ---	1,22
A	WO 96 32380 A (PHARMACIA S.P.A.) 17 October 1996 (1996-10-17) cited in the application claims -----	1,22

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 April 2000

Date of mailing of the international search report

02/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5816 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. J. Appl. No.

PCT/EP 99/09673

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640116 A	19-12-1996	US 5880141 A	09-03-1999
		AU 706597 B	17-06-1999
		AU 6044196 A	30-12-1996
		BR 9606410 A	30-12-1997
		CA 2192797 A	19-12-1996
		DE 29623744 U	30-09-1999
		EP 0769947 A	02-05-1997
		EP 0934931 A	11-08-1999
		HU 9701694 A	28-06-1999
		JP 2000026412 A	25-01-2000
		JP 10504323 T	28-04-1998
		NO 965377 A	12-02-1997
		NZ 310109 A	28-01-1999
		US 5792783 A	11-08-1998
		US 5883116 A	16-03-1999
		US 5834504 A	10-11-1998
		US 5886020 A	23-03-1999
		US 5883113 A	16-03-1999
WO 9632380 A	17-10-1996	EP 0764152 A	26-03-1997
		JP 10501821 T	17-02-1998
		US 5849710 A	15-12-1998

